

## SHORT REPORT

# Changes in outdoor air pollution due to COVID-19 lockdowns differ by pollutant: evidence from Scotland

Ruaraidh Dobson , Sean Semple 

Institute of Social Marketing and Health, University of Stirling, Stirling, UK

## Correspondence to

Dr Ruaraidh Dobson, Institute for Social Marketing and Health, University of Stirling, Stirling FK9 4LA, UK; r.p.dobson@stir.ac.uk

Received 29 April 2020

Revised 26 August 2020

Accepted 29 August 2020

Published Online First

7 September 2020

## ABSTRACT

**Objectives** To examine the impact of COVID-19 lockdown restrictions in March/April 2020 on concentrations of nitrogen dioxide (NO<sub>2</sub>) and ambient fine particulate matter (PM<sub>2.5</sub>) air pollution measured at roadside monitors across Scotland by comparing data with previous years.

**Methods** Publicly available data of PM<sub>2.5</sub> concentrations from reference monitoring systems at sites across Scotland were extracted for the 31-day period immediately following the imposition of lockdown rules on 23 March 2020. Similar data for 2017, 2018 and 2019 were gathered for comparison. Mean period values were calculated from the hourly data and logged values compared using pairwise t-tests. Weather effects were corrected using meteorological normalisation.

**Results** NO<sub>2</sub> concentrations were significantly lower in the 2020 lockdown period than in the previous 3 years ( $p < 0.001$ ). Mean outdoor PM<sub>2.5</sub> concentrations in 2020 were much lower than during the same period in 2019 ( $p < 0.001$ ). However, despite UK motor vehicle journeys reducing by 65%, concentrations in 2020 were within 1 µg/m<sup>3</sup> of those measured in 2017 ( $p = 0.66$ ) and 2018 ( $p < 0.001$ ), suggesting that traffic-related emissions may not explain variability of PM<sub>2.5</sub> in outdoor air in Scotland.

**Conclusions** The impact of reductions in motor vehicle journeys during COVID-19 lockdown restrictions may not have reduced ambient PM<sub>2.5</sub> concentrations in some countries. There is also a need for work to better understand how movement restrictions may have impacted personal exposure to air pollutants generated within indoor environments.

## INTRODUCTION

In the wake of the COVID-19 pandemic, many countries introduced wide-ranging restrictions on individual movement and gathering, known as 'lockdowns' or 'stay-at-home orders'. In the UK, a lockdown was introduced at 20:30 on 23 March 2020.

These new regulations led to substantial falls in road traffic with UK data suggesting motor vehicle journeys reduced by around 65% between 16 March and 28 April 2020.<sup>1</sup> The result of movement restrictions and reduced traffic volumes has been widely reported in the media (and some scientific studies) to have resulted in improved air quality and lower concentrations of common pollutants, such as fine particulate matter (PM<sub>2.5</sub>) and nitrogen dioxide (NO<sub>2</sub>).<sup>2,3</sup> It has been suggested that this will result in positive health effects, due to lowered exposure to air pollution, and even that the net effect of the

## Key messages

### What is already known about this subject?

- ▶ Road traffic has been significantly reduced in countries adopting lockdowns due to COVID-19. Research has shown that this has led to reductions in outdoor air pollution in some locations.

### What are the new findings?

- ▶ Nitrogen dioxide concentrations declined in Scotland following the lockdown, but fine particulate matter did not despite the fall in vehicle use.

### How might this impact on policy or clinical practice in the foreseeable future?

- ▶ Policymakers should take care not to overestimate improvements in outdoor air quality following COVID-19 lockdowns and should consider the impact of indoor air pollution on personal exposure during these periods.

pandemic will be to improve health (due to the adverse health effects of exposure to air pollution, particularly PM<sub>2.5</sub>).<sup>4</sup>

Analyses of this kind assume that road traffic-related PM<sub>2.5</sub> is a significant source of personal exposure to fine particles. This may not be true in all locations. Scotland's relatively low ambient PM<sub>2.5</sub> may be related more closely to natural and non-traffic sources and may not therefore have fallen following the introductions of the lockdown measures. If PM<sub>2.5</sub> in outdoor air has not declined, it is possible that net exposure to PM<sub>2.5</sub> will increase, as people spend more time in their homes where generation of fine particles from activities such as cooking and smoking may produce high concentrations within enclosed and poorly ventilated spaces.<sup>5</sup> NO<sub>2</sub> is specifically associated with vehicle exhaust emissions<sup>6</sup> and so provides a measure of relative traffic for use in this analysis.

## METHODS

Scottish local authorities maintain a network of automatic monitoring stations for PM<sub>2.5</sub> and other pollutants. The PM<sub>2.5</sub> monitors in use comprise gravimetric monitors (using tapered element oscillating microbalances, TEOMs) and high-precision optical monitors (optical aerosol spectrometers).



© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Dobson R, Semple S. *Occup Environ Med* 2020;**77**:798–800.

**Table 1** Geometric mean PM<sub>2.5</sub> and NO<sub>2</sub> in Scotland 24 March–23 April in years 2017–2020, including both observed and normalised data

Pollutant	2017 period geometric mean concentration (µg/m <sup>3</sup> )	2018 period geometric mean concentration (µg/m <sup>3</sup> )	2019 period geometric mean concentration (µg/m <sup>3</sup> )	2020 period geometric mean concentration (µg/m <sup>3</sup> )
PM <sub>2.5</sub> (observed)	6.7	7.4	12.8	6.6
PM <sub>2.5</sub> (normalised)	6.9	6.4	7.6	6.1
NO <sub>2</sub> (observed)	21.9	23.7	22.4	9.9
NO <sub>2</sub> (normalised)	25.8	25.4	24.4	15.1

NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, fine particulate matter.

These monitors report PM<sub>2.5</sub> measurements hourly and data are made publicly available on the internet.

To examine the effect of the lockdown on Scotland's air, PM<sub>2.5</sub> and NO<sub>2</sub> data were extracted from the monitor network for the period from 24 March to 23 April in 2017, 2018, 2019 and 2020. Data from 2020 have only been provisionally validated by the Scottish Government. Data were downloaded using the openair R package.

To simulate the removal of weather effects on pollutant concentrations, meteorological normalisation using the random forest machine learning algorithm<sup>7</sup> was conducted using the rmweather R package. Individual models were calculated for both PM<sub>2.5</sub> and NO<sub>2</sub> at monitoring sites around Scotland. Models were based on daily mean pollutant concentrations and incorporated wind speed, wind direction, atmospheric pressure, air temperature and relative humidity at the nearest available weather station (downloaded using the worldmet R package). Models used 64 trees and 100 samples.


Arithmetic mean concentrations were calculated for each of 70 PM<sub>2.5</sub> monitoring stations and 89 NO<sub>2</sub> monitoring stations over this period in each year. Geometric means of these values were calculated for each local authority area where monitoring took place and for Scotland overall in each year.

To determine statistical significance in differences in 2020 PM<sub>2.5</sub> and NO<sub>2</sub> values for this month versus each other year, both observed and normalised data were log-transformed and compared using a pairwise t-test. Statistical analysis was performed in R V4.0.2.<sup>8</sup>

## RESULTS

Across Scotland's air pollution monitoring network, observed and normalised NO<sub>2</sub> concentrations remained close to constant in 2017, 2018 and 2019 but fell substantially in 2020 (pairwise t-test  $p < 0.001$  for all years) (table 1).

By contrast, the observed geometric mean PM<sub>2.5</sub> concentration over the lockdown period in 2020 was 6.6 µg/m<sup>3</sup>, very similar to



**Figure 1** Observed geometric mean PM<sub>2.5</sub> and NO<sub>2</sub> by local government divisions (council areas) in Scotland 24 March–23 April in years 2017–2020. Note that some local authorities have NO<sub>2</sub> monitors but not PM<sub>2.5</sub> monitors. NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, fine particulate matter.

the mean concentration over the same period in 2017 (6.7 µg/m<sup>3</sup>, pairwise t-test  $p = 0.66$ ). The 2020 value showed a modest decrease ( $-0.8$  µg/m<sup>3</sup>) in comparison with 2018 (7.4 µg/m<sup>3</sup>,  $p < 0.001$ ) but was substantially lower than the markedly high concentrations measured in 2019 (12.8 µg/m<sup>3</sup>,  $p < 0.001$ ). Geometric means of normalised data showed the same pattern, with the 2019 mean higher than the other 3 years (pairwise t-test  $p < 0.001$  for all comparisons) (table 1).

Year 2019 was a visible outlier in observed data across all local authority areas where PM<sub>2.5</sub> monitoring was conducted (figure 1). This is likely due to a sustained meteorological event that brought fine particulate dust from the Saharan desert to the UK atmosphere beginning on 15 April 2019 and persisting through the end of the analysis period on 23 April.<sup>9</sup> Removing that period from the 2019 analysis reduces the mean observed value to 7.8 µg/m<sup>3</sup>, similar to overall values from the three other years in this analysis.

## DISCUSSION

The lockdown period has provided a natural experiment to examine the potential impact of reducing car journeys on air quality in Scotland. The NO<sub>2</sub> data suggest that car journeys have declined substantially during the lockdown compared with the same period in the previous 3 years. This may lead to significant health benefits, both from reduced exposure to harmful NO<sub>2</sub> and in reduced rates of traffic accidents and pedestrian collisions.

However, our results suggest that the decline in vehicle-related NO<sub>2</sub> has not coincided with significantly reduced PM<sub>2.5</sub> concentrations. The health risks of exposure to PM<sub>2.5</sub> are extremely well established, including cardiovascular disease, pulmonary illness and stroke. This research has established that reducing the number of vehicles on the road would not be an effective measure to reduce exposure to this pollutant in Scotland and consequently would not affect incidence of these illnesses.

Our analysis is limited by the data available from the monitoring network. Seven Scottish local authority areas have no NO<sub>2</sub> monitors, while nine have no PM<sub>2.5</sub> monitors, so these data do not cover the entirety of Scotland. Data from 2020 have been provisionally validated by the Scottish Government; while they have undergone screening to identify faulty or suspect data, they have not been ratified following detailed manual review. The later discovery of a fault or error associated with a monitor could change these results retroactively (if, for instance, a new calibration factor were applied). This is unlikely; in summer 2018, three faults were identified in particle monitors across the Scotland-wide network.<sup>10</sup> The use of data from a wide range of sources (70 PM<sub>2.5</sub> monitors and 90 NO<sub>2</sub> monitors) would limit the impact of a change to an individual monitor.

We have attributed the fall in normalised NO<sub>2</sub> concentrations in 2020 to the lockdown, but underlying effects, including a move towards less-polluting fuels and vehicles, could have contributed to this decline (though likely gradually over a period of years).

We believe these results have important policy and health implications in terms of the use of lockdowns to control future epidemics of infectious disease, and in considering how best to tackle outdoor air pollution in different countries in the future. Lockdowns are

intended to result in people spending more time in their homes. This could increase population exposure to indoor air pollution such as cooking fumes and secondhand tobacco smoke (a particular concern given the high concentrations of PM<sub>2.5</sub> that can be generated by smoking indoors). Previous work suggests that living with a smoker can increase a person's daily dose of PM<sub>2.5</sub> by over 80%.<sup>11</sup>

In countries like Scotland where it appears that the lockdown has not led to reductions in outdoor fine particulate matter pollution, it is possible that personal exposure to PM<sub>2.5</sub> may actually have increased rather than declined due to higher concentrations from indoor sources of particulate within the home setting. This could increase adverse health effects overall and also health inequalities—lower income people are more likely to smoke and to smoke indoors<sup>12</sup> and are likely to have smaller homes leading to higher PM<sub>2.5</sub> concentrations from individual sources, due to smaller room volumes. If the severity of COVID-19 is related to air pollution exposure (as has been suggested in ref 13), increased exposure to PM<sub>2.5</sub> could potentially increase the death toll of that disease. Careful and balanced consideration of both outdoor and indoor sources of PM<sub>2.5</sub> is essential to tackling the health harm of air pollution effectively and equitably.

**Contributors** Both authors conceived of the idea for the study. Both authors designed the study. RD conducted data analysis and drafted the manuscript, which SS critically reviewed. SS supervised the project.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. Data used in this study are available from the Scottish Government's air quality repository ([www.scottishairquality.scot](http://www.scottishairquality.scot)).

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise

determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

#### ORCID iDs

Ruaraidh Dobson <http://orcid.org/0000-0001-8136-8373>

Sean Semple <http://orcid.org/0000-0002-0462-7295>

#### REFERENCES

- 1 Prime Minister's Office. Slides and datasets to accompany coronavirus press conference, 2020. Available: <https://www.gov.uk/government/publications/slides-and-datasets-to-accompany-coronavirus-press-conference-28-april-2020> [Accessed 29 Apr 2020].
- 2 Duteil F, Baker JS, Navel V. COVID-19 as a factor influencing air pollution? *Environ Pollut* 2020;263:114466.
- 3 Cadotte M. Early evidence that COVID-19 government policies reduce urban air pollution.
- 4 Polichetti G, Cocco S, Spinali A, et al. Effects of particulate matter (PM(10), PM(2.5) and PM(1)) on the cardiovascular system. *Toxicology* 2009;261:1–8.
- 5 Klepeis NE, Bellettiere J, Hughes SC, et al. Fine particles in homes of predominantly low-income families with children and smokers: key physical and behavioral determinants to inform indoor-air-quality interventions. *PLoS One* 2017;12:e0177718.
- 6 Costa S, Ferreira J, Silveira C, et al. Integrating health on air quality assessment—review report on health risks of two major European outdoor air pollutants: PM and NO<sub>2</sub>. *J Toxicol Environ Heal Part B* 2014;17:307–40.
- 7 Grange SK, Carslaw DC. Using meteorological normalisation to detect interventions in air quality time series. *Sci Total Environ* 2019;653:578–88.
- 8 R Core Team. R: a language and environment for statistical computing., 2019. Available: <https://www.r-project.org/>
- 9 News BBC. Saharan dust: why your car is covered in red dirt, 2019. Available: <https://www.bbc.co.uk/news/uk-scotland-48037362> [Accessed 29 Apr 2020].
- 10 Ricardo Energy and Environment. *Scottish air quality database annual report 2018*, 2019.
- 11 Semple S, Apsley A, Azmina Ibrahim T, et al. Fine particulate matter concentrations in smoking households: just how much secondhand smoke do you breathe in if you live with a smoker who smokes indoors? *Tob Control* 2015;24:e205–11.
- 12 Scottish Government Community Health & Social Care Directorate. *Scottish health survey 2018: key findings*, 2019. Available: <https://www.gov.scot/publications/scottish-health-survey-2018-summary-key-findings/> [Accessed 29 Apr 2020].
- 13 Conticini E, Frediani B, Caro D. Can atmospheric pollution be considered a co-factor in extremely high level of SARS-CoV-2 lethality in northern Italy? *Environ Pollut* 2020;261:114465.