Livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) colonisation and infection among livestock workers and veterinarians: a systematic review and meta-analysis

Chen Chen ,¹ Felicia Wu^{1,2}

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2020-106418>).

¹Food Science and Human Nutrition, Michigan State University, East Lansing, Michigan, USA
²Agricultural, Food, and Resource Economics, Michigan State University, East Lansing, MI, United States

Correspondence to

Dr Felicia Wu, Food Science and Human Nutrition, Michigan State University, East Lansing, MI 48824, USA; fwu@msu.edu

Received 13 January 2020
Revised 21 September 2020
Accepted 1 October 2020
Published Online First
23 October 2020

ABSTRACT

Objectives Methicillin-resistant *Staphylococcus aureus* (MRSA) is an increasing public health concern worldwide. The objective of this study was to calculate a summary odds ratio (OR) of livestock-associated MRSA colonisation and infection in humans, and to determine specific risk factors in livestock production contributing to MRSA colonisation.

Methods We screened PubMed and Embase for studies published from 2005 to 2019 inclusive, reporting livestock-associated (LA)-MRSA colonisation and infection among livestock workers/veterinarians, their families, and community members not regularly exposed to livestock. The primary outcome of interest was the OR of LA-MRSA colonisation comparing exposed and control groups. Quality was assessed according to the Newcastle-Ottawa quality assessment scale. A meta-analysis using a random-effects model was conducted to calculate a pooled OR. The heterogeneity in the meta-analysis was assessed using the I^2 method, and publication bias was evaluated using funnel plots.

Results A total of 3490 studies were identified by the search, with 37 studies including 53 matched exposed-control groups and 14 038 participants eligible for the meta-analysis. The pooled OR for LA-MRSA among livestock workers and veterinarians is 9.80 (95% CI 6.89 to 13.95; $p=0.000$; $I^2=73.4$), with no significant publication bias (Egger's $p=0.66$). The OR for swine workers was highest at 15.41 (95% CI 9.24 to 25.69), followed by cattle workers (11.62, 95% CI 4.60 to 29.36), veterinarians (7.63, 95% CI 3.10 to 18.74), horse workers (7.45, 95% CI 2.39 to 23.25), livestock workers (5.86, 95% CI 1.14 to 30.16), poultry workers (5.70, 95% CI 1.70 to 19.11), and industrial slaughterhouse workers (4.69, 95% CI 1.10 to 20.0).

Conclusions Livestock workers, particularly swine farmers, are at significantly higher risk for LA-MRSA colonisation and subsequent infection. These results support the need for preventive practices to reduce LA-MRSA risk among those who handle and treat livestock.

Trial registration number CRD42019120403.

INTRODUCTION

Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a major cause of antibiotic resistant infections and a broader public health concern worldwide. As one of the most common pathogens in hospital settings, healthcare-associated MRSA (HA-MRSA) is a major cause of skin and soft tissue infections (SSTIs) and severe diseases such

Key messages

What is already known about this subject?

► Methicillin-resistant *Staphylococcus aureus* (MRSA) carriage is prevalent in livestock farmers, veterinarians, slaughterhouse workers, and butchers. However, whether the MRSA strains were livestock associated (LA) was less well-understood.

What are the new findings?

► Livestock workers, particularly swine farmers (with an OR of 15.41), are at significantly higher risk for LA-MRSA colonisation and subsequent infection compared with control populations.

How might this impact on policy or clinical practice in the foreseeable future?

► Preventive practices to reduce LA-MRSA risk are recommended among those who handle and treat livestock. In addition, coordinated strategies along livestock supply chains are recommended in farms, especially from regions with high risk of LA-MRSA colonisation.

as pneumonia and sepsis.¹ Twenty-nine per cent of MRSA carriers develop MRSA infection, which increases length of stay in hospitals, additional in-hospital costs and causes thousands of deaths worldwide.² In the USA, MRSA infections killed over 11 000 hospitalised patients in 2011.³

Since the 1990s, epidemiological characteristics of MRSA have changed greatly. New MRSA lineages have emerged, acquiring different types of the resistance genes responsible for infections occurring in the community (CA-MRSA, community-acquired MRSA).⁴ MRSA has also become widely distributed among livestock due to routine antibiotic use in livestock farming.⁵ Livestock-associated (LA)-MRSA strains are different from human MRSA strains, with a higher capacity of establishing and transmitting within livestock herds and potential for spreading and causing infections in humans.⁶ International trade of livestock has further intensified transmission of LA-MRSA between animals and humans.⁵ As global meat production and consumption have increased rapidly, farmers are facing pressures to produce livestock more efficiently to meet this demand. Often this involves administration of antibiotics worldwide, which can increase the risk of MRSA associated with livestock production.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chen C, Wu F. *Occup Environ Med* 2021;**78**:530–540.

LA-MRSA strains can be characterised by multilocus sequence typing, consist of *S. aureus* protein A (*spa*) and have a transmissible genetic element called staphylococcal cassette chromosome *mec*, which carries an element for beta-lactam antibiotic resistance.⁷ An increasing body of evidence demonstrates that livestock workers are a high-risk population for LA-MRSA carriage due to regular close contact with animals, especially in farm-intensive regions. LA-MRSA has emerged in swine production globally. The prevalence of LA-MRSA colonisation varies widely among livestock workers throughout the world, with reported rate in pig workers of 24%–86% in Europe, 25%–45% in North America, and 6%–19% in Asia.^{8–11} The epidemiology of LA-MRSA strains differs in different geographic regions. In Europe, MRSA ST398 is the predominant clone complex found in livestock workers or veterinarians. Other clones such as ST1 (the USA), ST5 (the USA and Ireland), ST8 (the USA, Denmark and Switzerland), and ST22 (Ireland) were also identified at lower frequencies in livestock operations. In Asian countries, the presence of various clones of LA-MRSA such as ST9, ST63 and ST2359 were reported in livestock workers.^{12–14}

LA-MRSA isolates have a strong adherence capacity to human keratinocytes, and humans working closely with livestock harbouring LA-MRSA isolates may become colonised.¹⁵ LA-MRSA can cause severe SSTIs, which mainly occur in persons with occupational exposure to livestock. LA-MRSA is capable of being established in hospital settings via colonised or infected patients, which can lead to nosocomial infections, although the incidence of these infections is rare.⁷

A prior systematic review summarised the prevalence of MRSA carriage in livestock workers: farmers (18.2%), veterinarians (9.4%), slaughterhouse workers (2.6%) and butchers (5.7%).¹⁶ However, whether the MRSA strains were human-associated or livestock-associated was not distinguished, and the extreme heterogeneity of the results ($I^2=96.9\%$) called into question whether a pooled prevalence estimate such as they provided offered a meaningful statistic. In another study, a 760-fold higher risk of MRSA colonisation was found among a group of regional pig farmers compared with the general population.¹⁷

What has already been uncovered about the risk of livestock-acquired MRSA comes from individual studies examining, mostly, the impact of one type of occupation (eg, swine producer, cattle producer and veterinarian) on the risk of LA-MRSA colonisation. These studies range across multiple different species and multiple different occupations, providing a range of odds ratios (ORs) and relative risks when comparing the exposed population to a control population.

The objective of the current study is to calculate a summary OR of LA-MRSA colonisation and infection in humans who are occupationally exposed to livestock and to determine specific risk factors in livestock production contributing to MRSA colonisation. Our unique contribution in this manuscript is to bring together these studies in a systematic review and meta-analysis: to examine and quantify overall risk of occupational LA-MRSA from multiple different nations worldwide and to pool the data from existing studies on each type of occupation to determine that livestock-related occupations result in the highest LA-MRSA risk.

METHODS

Data sources and search strategy

We performed a systematic review and meta-analysis of the risk of livestock contact in increasing the incidence of MRSA colonisation in occupational workers and veterinarians. A literature search was conducted on PubMed and Embase databases from 31 October to 21 October 2019. The search terms included combinations of: (“methicillin-resistant *Staphylococcus aureus*”

OR “MRSA”) AND (“livestock” OR “pig” OR “swine” OR “hog” OR “cattle” OR “poultry” OR “farmer” OR “worker”). Retrieved publications were reviewed by examining titles and abstracts, and all potentially relevant articles were subsequently screened as full text. In addition, the search was extended by examining the reference lists of retrieved publications to find more relevant studies linking MRSA carriage with livestock farming or care. No study type or language restriction was imposed. The meta-analyses were conducted according to Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines (research checklist). The leading author (CC) screened the search results and identified relevant papers by reviewing retrieved studies. The quality assessment of included studies was conducted by both authors together. Any discrepancies on inclusion were resolved by joint reevaluation of the original full text by both authors. Studies reporting the occurrence of LA-MRSA colonisation among workers/veterinarians and control populations were included in the analysis. The microbiological method based on strain typing was a criterion for presence of LA-MRSA. Studies without control groups, or that did not differentiate between LA-MRSA and other types of *Staphylococcus aureus*, were excluded. This systematic review is registered with PROSPERO.

Data extraction

The primary outcome of interest was LA-MRSA colonisation among study participants who had close contact with livestock, determined by LA-MRSA strain isolation from nasal, oropharyngeal, or axillary samples. The participants in the eligible studies were divided into exposed and control groups according to their occupational contact with livestock. The exposed group includes the following populations who have direct contact with animals: livestock farmers and veterinarians (including veterinary students) with occupational exposure to various animal species, and slaughterhouse workers including butchers, industrial hog operation and industrial livestock operation workers. The control populations are those who have close proximity to colonised animals but no or minimal direct contact with them, including household and community members of farmers, employees working on farms (eg, salesmen and administrative personnel), antibiotic-free livestock operation workers, and employees not working on farms or treating livestock. These categories allowed the examination of OR of LA-MRSA colonisation among occupations with different levels of exposure to different livestock species.

The following data were extracted from each study: authors, year of publication, study location, study period and duration, study design type, population size, participants' gender and age (range), type of occupations, strain type and genotype identified, and LA-MRSA colonisation rate and infections among exposed and control participants. In addition, the details of the screening including sampling sites (nasal, oropharyngeal or axillary swabs), MRSA isolation method (culture or PCR), genotyping of isolates, carriage status definition and timeframe of the screening process were retrieved from the studies. Regarding detection methods of microbial isolates, two studies applied a culture method (enrichment broth, sheep blood and chromogenic medium) to screen for carriage of MRSA isolates, which is less sensitive than PCR assays to determine the genotype of MRSA. They were excluded from the analysis. When needed, authors were contacted via email to request additional information.

To ensure that only LA-MRSA strains are included in the meta-analysis, numbers of carriers with *spa* types not of animal origin

Figure 1 Study selection flow chart. LA-MRSA, livestock-associated methicillin-resistant *Staphylococcus aureus*; MRSA, methicillin-resistant *Staphylococcus aureus*.

were subtracted from the LA-MRSA positive persons.^{18–20} The secondary outcome of interest was the OR of developing MRSA infections among exposed and control populations. In addition, an index year of each eligible study was determined to model the time trend of LA-MRSA colonisation among participants.

Study quality assessment

Quality assessment of included studies was conducted by both authors according to the Newcastle-Ottawa Scale, a star-based scoring system. Studies that received at least five stars were considered of adequate quality for eligibility.

Data analysis

The OR is a comparison of the odds of an event after exposure to a risk factor (colonisation of LA-MRSA in this analysis) with the odds of that event in a control situation. For each study, matched exposed-control groups were identified, from which ORs and 95% CIs could be calculated between the different exposed and control groups. A random-effects model meta-analysis was conducted to calculate the pooled OR and the associated 95% CI from the eligible studies. Heterogeneity among the selected studies was evaluated using the Cochran's Q value calculated from the Mantel-Haenszel method and I^2 statistic. Sensitivity analyses were conducted by removing each study in turn, and the remaining studies were analysed to determine if the results were significantly impacted by a particular study. The effect of the index year of the study as a covariate on OR of LA-MRSA colonisation was explored through meta-regression analysis for each continent. P values for OR comparisons were calculated using the Fisher's exact test. Bias in meta-analysis was evaluated

by asymmetry in funnel plot and related statistical analyses. All statistical analyses were conducted in Comprehensive Meta-Analysis Software V2.0.

RESULTS

Our literature search retrieved 3490 studies based on our search terms (figure 1). After applying the eligibility criteria described above at the title and abstract level, 2363 studies were excluded. We then retrieved and read the full texts of the remaining 218 studies, from which 35 were selected for sufficient data for our systematic review and meta-analysis. Two of the contacted authors were able to provide information included in the analysis. Two additional articles were identified from search of the references of the 35 included studies. Finally, 37 studies were included in this systematic review and meta-analysis.^{12–14 18 21–53} Six studies were of six stars, 32 received five stars (online supplemental material table S1).

An overview of the eligible studies is listed in table 1, including the year(s), location and study type; number and occupation of participants and their gender and age; sample type and measurement of colonisation; *spa* type; number of LA-MRSA carriers in exposed and control populations; and the number of MRSA infection cases. The 37 eligible studies on LA-MRSA colonisation and occupational livestock exposure consist of 26 cross-sectional studies, 5 prospective cohort studies, 3 case-control studies and 3 longitudinal studies published between 2005 and 2019 from nine countries, comprising 53 exposed-control groups and 14 038 participants. Twenty-one studies assessed the risk of MRSA carriage in occupational pig workers, 10 in veterinarians, 7 in cattle workers and 5 in poultry workers.

The overall pooled OR of LA-MRSA carriage among exposed populations based on the random-effects model was 9.80 (95% CI 6.89 to 13.95; $p=0.000$; $I^2=73.4$) compared with control populations, among 53 exposed-control subgroups, indicating that livestock contact is a significant risk factor for LA-MRSA colonisation in humans. Stratifying the studies by geographical regions, the pooled OR from Asian countries (farm workers vs non-farm workers) was 4.03 (95% CI 1.44 to 11.29; $p=0.15$; $I^2=46.9$). Among European countries, the pooled OR was 9.56 (95% CI 6.90 to 13.25; $p=0.000$; $I^2=76.5$). A relatively lower OR (4.69, 95% CI 1.10 to 20.05; $p=0.04$; $I^2=0.000$) was found in the USA. Figure 2 shows classifications and population size of subgroups according to their occupational contact with livestock farming and the forest plots for the pooled OR and 95% CI of LA-MRSA colonisation among all exposed subgroups.

When stratifying occupations of the study participants by different animal species, the highest OR was seen in pig workers (15.41; 95% CI 9.24 to 25.69; $p=0.000$; $I^2=68.7$), followed by cattle workers (11.62, 95% CI 4.60 to 29.36; $p=0.000$; $I^2=79.0$), veterinarians (7.63, 95% CI 3.10 to 18.74; $p=0.000$; $I^2=77.9$), livestock workers (5.86, 95% CI 1.14 to 30.16; $p=0.23$; $I^2=32.6$), poultry workers (5.70, 95% CI 1.70 to 19.11; $p=0.006$; $I^2=72.3$) and industrial operation workers (4.69, 95% CI 1.10 to 20.0; $p=0.04$; $I^2=0.000$). Horse workers were 7.45 times (95% CI 2.39 to 23.25; $p=0.17$; $I^2=47.6$) more likely to be colonised by LA-MRSA compared with control participants based on two relevant studies. One study reporting risk of LA-MRSA colonisation among patients with sheep contact did not show a different risk (1.0, 95% CI 0.14 to 7.24) than control population.

Medium to substantial heterogeneity ($I^2 \geq 50$) was observed in the subgroups of cattle farmers, veterinarians and poultry workers. The sensitivity analyses show that excluding the most

Table 1 Summary of studies included in the meta-analysis (n=37)

	Country	Study type	Period	Sex*	Age	Exposed population	Control population	Samples	Main Spa types of LA-MRSA	Non-LA-MRSA colonisation	Infection
Angen <i>et al</i> ²¹	Denmark	One-week longitudinal study.	2016	27/7	–	48 volunteers visiting a swine farm.	46 passive observers.	Nasal and throat swabs.	LA-MRSA CC398 (t011).	–	–
Bisdorff <i>et al</i> ¹⁸	Germany	Cross-sectional study.	2009–2010	Non-OLC: 913/742; OLC: 581/32.	Non-OLC: 42.2; OLC: 42.7.	1655 OLC group.	190 non-OLC group.	Nasal swabs.	ST398 (t034 and t011).	7 out of 25 non-OLC carried HA-MRSA.	–
Bos <i>et al</i> ²²	The Netherlands	Longitudinal study	–	291/371	30.7 (0–83)	182 pig farmers and 67 cattle farmers.	286 family members of pig farmers and 127 of cattle farmers.	Nasal swabs.	ST398	–	–
Cuny <i>et al</i> ²³	Germany	Cross-sectional study.	September 2007–January 2009	–	–	113 pig farmers; 49 veterinarians.	116 non-exposed family members of pig farmers; 44 non-exposed family members of veterinarians.	Nasal swabs.	ST398 (t011, t034, t2974 and t108).	–	–
Denis <i>et al</i> ²⁴	Belgium	Cross-sectional.	April–July 2007	–	–	75 pig farmers.	27 persons without pig contact.	Nasal swabs and wound.	ST398 (t011, t034 and t567).	–	One LA-MRSA carrier had hand lesion infection.
Fang <i>et al</i> ²	Taiwan	Cross-sectional study.	June–October 2012	16/84	–	52 pig farm workers.	32 auction market employers and 16 regular visitors.	Nasal swabs.	ST9 (t899 and t1939).	Three farmers carried CA-MRSA.	–
Geenen <i>et al</i> ²⁷	The Netherlands	Cross-sectional study.	July 2010–May 2011.	–	Mean: 36.	47 farmers.	89 family members and 9 employees.	Nasal swabs.	ST398 (t011, t034 and t0108).	–	–
Gracia-Graells <i>et al</i> ^{26,4}	Belgium, Denmark and The Netherlands	6 months longitudinal study.	2009–2010	–	–	15 pig farmers.	45 household members.	Nasal swabs.	ST398 (t011, t034, t0108 and t1451).	–	–
Gracia-Graells <i>et al</i> ²⁵	Belgium	Cross-sectional study, Belgium.	February–April 2010	50/96	45.1	105 veterinarians working with livestock.	41 veterinarians not working with livestock.	Nasal swabs.	ST398 (t011, t034, t1451 and so on).	–	–
Gracia-Graells <i>et al</i> ²⁵	Denmark	Cross-sectional study, Denmark.	February–October 2010	–	–	97 veterinarians working with livestock.	46 veterinarians not working with livestock.	Nasal swabs.	ST398 (t011, t034, t1451 and so on).	–	–
Graveland <i>et al</i> ²⁸	The Netherlands	Cross-sectional study.	October 2007–March 2008	177/213	0–85	97 calf farmers.	259 family members and 34 employees.	Nasal swabs.	ST398 except four (t002, t015, t084 and t166).	1 HA-MRSA and 3 CA-MRSA carriers.	–
Hatcher <i>et al</i> ²⁹	USA	Cross-sectional study.	2014	Adult: 252/148; children: 187/213.	–	198 IHO worker–child household pairs.	202 community referent adult–child household pairs.	Nasal swabs.	ST5 (t002), ST8 (t008 and t088).	–	–
Huber <i>et al</i> ³⁰	Switzerland	Conference participants.	March–September 2009	–	–	148 pig farmers and 133 veterinarians.	179 slaughterhouse employees.	Nasal swabs.	All CC398 (ST8 t064, ST398 t011 and t034).	–	–
Köck <i>et al</i> ³¹	Germany	Case–control study.	January 2005–December 2008.	Case group: 23/77; control: 38/62.	Case group: 48; control: 54.	100 patients with LA-MRSA.	100 patients with other than LA-MRSA.	Nasal swabs.	ST398 (t011, t034, t108 and t1451).	–	–
Lewis <i>et al</i> ³²	Denmark	Case–control study.	2004–2007	Case-patients: 13/8.	Median of case-patients 29 (8 months–80 years).	21 case patients with CC398 MRSA.	42 controls.	MRSA isolates from patients.	ST398 (t034, t108 and t1793).	–	10 case-patients had reported SSTI.
Moodley <i>et al</i> ³³	Denmark	Conference participants.	August 2006–February 2007	–	–	231 veterinarians, 72 veterinary students or nurses, 98 farmers.	301 unexposed persons.	Nasal swabs.	CC8 (t008), CC22 (t020 and t022), CC59 (t216), CC88 (t186) CC398 (t011 and t034).	Exposed: 6/9 were LA-MRSA carriers; unexposed: 1/2 was LA-MRSA carrier.	–
Mutters <i>et al</i> ³⁴	Germany	Cross-sectional study.	June–October 2012	–	–	63 occupationally exposed individuals.	126 patients.	Nasal and rectal swabs.	Farmers CC398; 4/126 control inpatients CC398.	–	–

continued

Table 1 continued

	Country	Study type	Period	Sex*	Age	Exposed population	Control population	Samples	Main <i>Spa</i> types of LA-MRSA	Non-LA-MRSA colonisation	Infection
Nadimpalli <i>et al</i> ³⁵	USA	Cross-sectional study.	October 2013–February 2014	94/89	Mean: 30.	103 IHO workers.	80 family members.	Nasal swabs.	CC398		6 IHO and 6 of 80 household members had SSTI.
Neyra <i>et al</i> ³⁶	USA	Cross-sectional study.	September–November 2011	196/140	15–82	162 hog slaughter workers.	63 household members of plant workers and 111 community residents.	Nasal swabs.	ST1 (LA-MRSA), ST5 (HA-MRSA), ST8, 1 of ST72 (CA-MRSA) and ST398.		
Oppliger <i>et al</i> ³⁷	Switzerland	Cross-sectional study.	June 2008–July 2009	–	–	75 pig farmers and veterinarians.	128 persons without animal contact.	Nasal swabs.	CC398.	–	
Pletinckx <i>et al</i> ³⁸	Belgium	Cross-sectional study.	July 2009–October 2010	–	–	10 farmers and 10 veterinarians.	13 family members (living in farms).	Nasal swabs.	ST398 (t011 t034 t567).		
Richter <i>et al</i> ³⁹	Germany	Cross-sectional study.	June–October 2009	–	–	39 people with frequent contact with turkeys.	20 persons being rarely or never in turkey houses.	Nasal swabs.	ST398 (t011 and t034).		
Rinsky <i>et al</i> ⁴⁰	USA	Cross-sectional study.	May–December 2011	ILO: 59/40; AFLO: 46/59.	–	80 ILO workers.	19 household members of ILO workers, 92 AFLO workers and their 13 household members.	Nasal swabs.	CC398		
Sahibzada <i>et al</i> ⁴¹	Australia	Cross-sectional study.	2015	12/40	–	37 piggy staff working in pig contact.	15 workers working in the agriculture or feedmill section.	Nasal swabs.	LA-MRSA ST398.	29 piggy staff carried ST93; two workers working in the agriculture or feedmill carried ST93.	
Schmithausen <i>et al</i> ⁴²	Germany	Cross-sectional study.	June–December 2012	–	–	79 pig farmers.	7 family members.	Nasal swabs.	ST398 (t011 and t034).		
van Cleef <i>et al</i> ^{44,45,†}	The Netherlands	Prospective cohort study.	2010–2011	103/68	Median: 16 (0–70).	110 pig farmers.	171 household members.	Nasal swabs.	MLVA complex (MC) 398.		
van den Broek <i>et al</i> ⁴⁶	The Netherlands	Cross-sectional study.	January–October 2007	107/125	0–86	50 farmers.	171 family members and 11 coworkers.	Anterior nasal swabs.	ST398 (t108, t011 and t567 and others).	–	
Van Cleef <i>et al</i> ⁴³	The Netherlands	Cross-sectional study.	2008	16/233	Mean: 43.	207 slaughterhouse workers, 13 veterinarians and auxiliaries.	29 administrative and technical personnel.	Nasal swabs.	ST398 (t011 and t108).		
Vandendriessche <i>et al</i> ⁴⁹	Belgium	Cross-sectional study.	August 2009–May 2011	–	–	149 farmers: pig (25), veal (45), dairy (22), beef (21) and broiler (36).	42 farmers and family members living on horticulture farms.	Nasal swabs.	CC398 (t011, t034 and t567).		
van Duijkeren <i>et al</i> ⁴⁷	The Netherlands	Cross-sectional study.	November 2013–September 2014	–	–	11 turkey farmers.	32 family members and employees.	Nasal swabs.	ST398 (t011)		
van Loo <i>et al</i> ⁴⁸	The Netherlands	Case–control study.	2003–2005	56/55	Case: 42.7; control: 47.2.	35 case patients with CC398.	76 control patients with non-CC398 MRSA.	NT-MRSA isolates from patients.	ST398 (t108, t011, t034 and t571).	–	
Verkade <i>et al</i> ⁵⁰	The Netherlands	A 1-year prospective cohort study.	2009–2011	–	18–65.	135 livestock veterinarians.	386 non-exposed household members.	Anterior nares or oropharynx swabs.	MC398.		
Walter <i>et al</i> ^{51,52}	Germany	Prospective cohort study.	2011–2014	–	Median of 45 for LA-MRSA participants.	1435 participants of veterinary congresses.		Nasal swabs.	CC398 (t011, t034 and t571).		
Wang <i>et al</i> ¹³	China	Cross-sectional study.	November 2013–November 2014	597/593	Workers: 39.3 controls: 36.4.	335 pig-related workers (including 178 slaughterhouse workers and 157 butchers).	855 control workers.	Nasal swabs.	Only 4 MRSA in workers and 3 in controls were LA.		

continued

Table 1 continued

	Country	Study type	Period	Sex*	Age	Exposed population	Control population	Samples	Main <i>Spa</i> types of LA-MRSA	Non-LA-MRSA colonisation	Infection
Wuif <i>et al</i> ²³	The Netherlands	Prospective surveillance study.	–	726/127	Mean: 36.9.	222 healthcare worker.	633 controls.	Throat and nasal swabs.	ST398 (1108).		
Ye <i>et al</i> ¹⁴	China	Cross-sectional study.	November 2013–November 2014	669/1191	15–60	682 occupational livestock workers (224 farm workers, 20 veterinarians, 194 slaughterhouse workers and 244 butchers).	1178 controls	Anterior nasal swabs.	CC9 (ST9, ST63 and ST2359).	Farmers: 18/48 were LA-MRSA; controls: 2/16 were LA-MRSA.	–

*The number of female and male participants.

†LA-MRSA carriers were those who were colonised by LA-MRSA persistently or intermittently.

#Tetracycline resistance was considered as the marker for livestock association among MRSA strains that showed resistance to cefoxitin.

AFLC, antibiotic free livestock operation; CA, community-associated; HA, healthcare associated; IHO, industrial hog operation; LA, livestock-associated; MLVA, multilocus variable-number-tandem-repeats analysis; OLC, occupational livestock contact; *Spa*, *Staphylococcus aureus* protein A; SSTI, skin and soft tissue infection.

heavily contributing study resulted in significantly reduced heterogeneity within these subgroups (figure 3).

The time trend plot (online supplemental material figure S1) on the basis of index year of the eligible studies indicated that a decreasing trend of LA-MRSA colonisation among livestock workers existed only in European countries (fixed effect, $p=0.046$), but not in North America ($p>0.5$) or Asia ($p>0.5$). Three of the selected studies recorded SSTI among the participants. The risk for MRSA infections based on these studies was 3.44 (95% CI 0.20 to 58.3; $p=0.015$; $I^2=76.1$) among occupational livestock workers. In hospital settings, infections with more serious symptoms and outbreaks caused by LA-MRSA among persons with occupational livestock contact have also been reported.

Publication bias was assessed by a funnel plot (online supplemental material figure S2). The associated statistical test (Egger's test $p=0.66$) of funnel plot asymmetry indicated that publication bias was not a serious limitation in our meta-analysis.

DISCUSSION

Our analysis indicates that livestock-exposed populations were 9.80-fold more likely to be colonised by LA-MRSA than the control population. In particular, pig farmers had the highest OR for LA-MRSA colonisation of 15.41. The differences in ORs for different livestock species may result from differences in basic facilities and operations across the species, or differences in transmissibility of LA-MRSA between each of these species and humans.

Our study has several limitations. The prevalence of LA-MRSA carriage in human population is associated with animal-to-human transmission, human-to-human transmission and airborne route.⁵⁴ Intensive contact with livestock allows LA-MRSA to persist in workers, and transmission events can occur on farms and beyond farms. Thus, the long-term persistence of the carriage is considered to play a key role in the transmission dynamics of LA-MRSA.⁵⁵ In this regard, a main limitation of our analysis is that the transmission dynamics of LA-MRSA was not taken into account when using the association between cause and effect in epidemiology. For the majority of the selected studies, whether the LA-MRSA carriage patterns were intermittent or persistent was not identified in studies that took cross-sectional nasal and/or throat samples; only six studies examined samples longitudinally at multiple time points in which occurrence of MRSA carriage over time is associated with the use of antibiotics and other determinants.^{21 26 45 50 51} New evidence suggests that colonisation of ST398 in humans is potentially more persistent; the transmission from farm workers to their household members occurs at a low but consistent percentage.^{2 23} Therefore, we assumed the carriage of LA-MRSA being in equilibrium, although this state is practically subject to change due to population dynamics and stochastic events in the transmission. Nonetheless, the weight of evidence shows that occupational livestock contact puts the workers at higher risk for LA-MRSA colonisation.⁵⁶

We should take into account that a single nasal or throat swab is not a proof of colonisation of LA-MRSA, which has been shown to be a relatively weak coloniser in humans, where carriage of MRSA can disappear in the absence of animal exposure. However, in the majority of the selected studies, nasal and/or throat swab samples were collected for the detection of LA-MRSA carriage, except two studies in which the strain isolates were collected from patients infected with LA-MRSA, which may overestimate the number of positive individuals and

Figure 2 Forest plots of the OR and 95% CI for studies of LA-MRSA colonisation among subgroups who have contact with different animal species. The squares indicate or estimate within each subgroup; the diamond at the bottom indicates the pooled OR estimate in the meta-analysis according to the random-effects model.

(a): Forest plots for OR of LA-MRSA colonisation among cattle workers excluding Bisdorff et al. (2012)

(b): Forest plots for OR of LA-MRSA colonisation among veterinarians excluding Walter et al. (2016; 2017)

(c): Forest plots for OR of LA-MRSA colonisation among poultry workers excluding Bisdorff et al. (2012)

Figure 3 Forest plots for one-study-removed meta-analysis of OR among cattle workers (A), poultry workers (B) and veterinarians (C). The plots show the OR estimate when each study is removed from the meta-analysis. The deviation from the full analysis indicates the sensitivity of the full analysis with respect to each study, indicating the weight a particular study contributes to the meta-analysis. LA-MRSA, livestock-associated methicillin-resistant *Staphylococcus aureus*.

therefore is another limitation of our study. Moreover, colonisation of ST398 in humans has the potential to be more persistent and a strong link exists between occupational animal contact and persistent LA-MRSA colonisation.^{2,44} Participants in the studies reviewed here generally have long-term repetitive contact with

livestock, which is a major risk factor for persistent LA-MRSA carriage in humans.

Because no associations of the risk of LA-MRSA colonisation were found for demographic information (dietary preferences, exposure to raw meat, smoking, contact sports and travel) and

medical history (exposure to healthcare facilities and antibiotic usage),^{12 23 26} these variables were not considered as risk factors in our meta-analysis. In intensive farming, some livestock workers take preventive measures during work to reduce the risk of bacterial infection. Unfortunately, these measures have not been broadly successful for preventing occupational acquisition of LA-MRSA. For example, the majority of industrial hog operators wear protective footwear, long sleeves, long trousers or coveralls when working in operation areas.³⁵ However, the OR (4.69) of LA-MRSA colonisation among this group was still significantly higher than control population. Moreover, implementing nationwide eradication programmes has proven difficult.⁵⁴ Although the majority of LA-MRSA colonisation cases in occupationally exposed persons may be asymptomatic, a quantitative association between LA-MRSA carriage and infection still needs to be investigated in the future research.

The time trend analysis shows a decreasing risk of LA-MRSA colonisation among livestock workers in European countries. This may be due to multiple different phenomena. Farming practices may be improving to reduce LA-MRSA transmission risk, including methods to keep animals healthy without the need for antibiotics; such as the use of vaccines, nutritional supplements, prebiotics, probiotics or synbiotics, and waste management strategies.⁵⁷ Veterinarians are more informed than ever about the risks of antimicrobial resistance; hence, they are careful in their own use of antibiotics with animals as well as the advice that they provide to livestock and poultry farmers. However, changes in the prevalence in control populations may also contribute to the change in risk of colonisation over time. We calculated the time trend of the risk difference of LA-MRSA carriage between populations with intensive animal contact and those in the control population and found that the risk difference remains stable over time ($p > 0.1$) based on a meta-regression analysis, which might suggest an equilibrium of prevalence of LA-MRSA between the exposed and control groups.

Most (28 out of 37) of the selected studies are originated from European countries, but the risk of LA-MRSA colonisation varied widely among countries. This may be because of different regulations or enforcement of regulations regarding animal operations (including antibiotic use) across countries or because of differences in climate (temperature and precipitation) that could affect MRSA transmission rates. Trading colonised animals between countries is a vehicle and considered as another major risk factor for transmission of LA-MRSA between humans and pigs.⁵⁸ In MRSA-free areas, livestock imports should be considered an important route of introduction of LA-MRSA into livestock populations.⁵⁹ The LA-MRSA strains in pig supplying farms from regions with high risk of LA-MRSA colonisation can highly affect the MRSA status of the recipient farms. In addition, associations have been observed between MRSA infections and high-density livestock production.⁷ Coordinated strategies along livestock supply chains are recommended to control and prevent LA-MRSA in livestock farming and trade.

The source of heterogeneity in the meta-analysis is also a limitation. Many studies providing the prevalence of LA-MRSA among livestock workers without a control population were excluded, prohibiting a calculation of an OR. Currently, there is lack of consensus on validated markers for detecting livestock-associated *S. aureus*, which may be a source of heterogeneity in our analysis. Absence of *scn*, positivity of *mecA* and tetracycline resistance are recognised as main indicators of livestock association among MRSA isolates.³⁵ However, these characteristics have also been found in other lineages. Some strains of LA-MRSA appear to be phylogenetically originated in humans

and evolved a variety of adaptations via transmission and spread among hosts.⁶⁰ As a result, some strains of LA-MRSA may be human associated. Second, no selected studies originated from Africa or South Asia, which influences the representativeness of our analysis but highlights the need for further research in these regions.

The risk of developing LA-MRSA colonisation among livestock workers and veterinarians is also affected by other risk factors: livestock density and the type of farm or stage of animal production (eg, LA-MRSA transmission risk was higher in sows than finishing pigs), antimicrobial use history, hospital admission, working on farms with MRSA-positive animals, herd size and sanitary conditions.^{23 46 56} These factors vary among individuals and different geographical regions, representing additional source of heterogeneity. However, these variables have not been measured in the majority of the selected studies. Our analysis summarised a combined estimate of the OR but was not able to identify the contributions of these risk factors to LA-MRSA colonisation among study participants.

The principal strength of our study is the analysis of the OR of LA-MRSA colonisation and infection among livestock workers and veterinarians as the main outcomes of the occupational contact with livestock. The pooled OR indicates a high risk and major burden of LA-MRSA colonisation and infection currently involved in livestock contact, which underscores the importance of this clone for global occupational health. Occupational exposure to pigs is the most important factor for LA-MRSA carriage, which underscores the importance of this clone for global occupational health. Future studies should focus on implementing interventions in livestock farming and operation and veterinary settings and determining the effectiveness of these interventions in reducing LA-MRSA colonisation among occupational workers and veterinarians.

CONCLUSION

The review shows that livestock workers, particularly swine farmers, are at significantly higher risk for LA-MRSA colonisation and subsequent infection. These results support the need for preventive practices to reduce LA-MRSA risk among those who handle and treat livestock. In addition, coordinated strategies along livestock supply chains are recommended in farms, especially from regions with high risk of LA-MRSA colonisation, to control and prevent LA-MRSA in livestock farming.

Contributors FW conceived and designed the study. CC did the literature review and data extraction. Both authors evaluated the studies. CC was involved in the data synthesis and statistical analysis. CC and FW contributed to the interpretation of the data and writing of the manuscript. Both authors agreed with its content and approved the final submitted version.

Funding This study was funded by USDA grant MICL02527 (FW).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iD

Chen Chen <http://orcid.org/0000-0003-2599-8332>

REFERENCES

- 1 Kluytmans J, Struelens M. Methicillin resistant *Staphylococcus aureus* in the hospital. *BMJ* 2009;338:b364.
- 2 Köck R, Loth B, Köksal M, et al. Persistence of nasal colonization with livestock-associated methicillin-resistant *Staphylococcus aureus* in pig farmers after holidays from pig exposure. *Appl Environ Microbiol* 2012;78:4046–7.
- 3 Dantes R, Mu Y, Belflower R, et al. National burden of invasive methicillin-resistant *Staphylococcus aureus* infections, United States, 2011. *JAMA Intern Med* 2013;173:1970–8.
- 4 WHO. *The work of who in the south-east Asia region, report of the regional director, 1 January-31 December 2016*, 2017.
- 5 Aires-de-Sousa M. Methicillin-resistant *Staphylococcus aureus* among animals: current overview. *Clin Microbiol Infect* 2017;23:373–80.
- 6 Elstrøm P, Grøntvedt CA, Gabrielsen C, et al. Livestock-Associated MRSA CC1 in Norway; introduction to pig farms, zoonotic transmission, and eradication. *Front Microbiol* 2019;10:139.
- 7 Cuny C, Wieler LH, Witte W. Livestock-associated MRSA: the impact on humans. *Antibiotics* 2015;4:521–43.
- 8 Khanna T, Friendship R, Dewey C, et al. Methicillin resistant *Staphylococcus aureus* colonization in pigs and pig farmers. *Vet Microbiol* 2008;128:298–303.
- 9 Smith TC, Male MJ, Harper AL, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS One* 2009;4:e4258.
- 10 Chuang Y-Y, Huang Y-C. Livestock-associated methicillin-resistant *Staphylococcus aureus* in Asia: an emerging issue? *Int J Antimicrob Agents* 2015;45:334–40.
- 11 Golding GR, Bryden L, Levett PN, et al. Livestock-associated methicillin-resistant *Staphylococcus aureus* sequence type 398 in humans, Canada. *Emerg Infect Dis* 2010;16:587–94.
- 12 Fang H-W, Chiang P-H, Huang Y-C. Livestock-associated methicillin-resistant *Staphylococcus aureus* ST9 in pigs and related personnel in Taiwan. *PLoS One* 2014;9:e88826.
- 13 Wang XL, Li L, Li SM, et al. Phenotypic and molecular characteristics of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* in slaughterhouse pig-related workers and control workers in Guangdong Province, China. *Epidemiol Infect* 2017;145:1843–51.
- 14 Ye X, Liu W, Fan Y, et al. Frequency-risk and duration-risk relations between occupational livestock contact and methicillin-resistant *Staphylococcus aureus* carriage among workers in Guangdong, China. *Am J Infect Control* 2015;43:676–81.
- 15 Hau SJ, Kellner S, Eberle KC, et al. Methicillin-resistant *Staphylococcus aureus* sequence type (ST) 5 isolates from health care and agricultural sources adhere equivalently to human keratinocytes. *Appl Environ Microbiol* 2018;84:e02073–17.
- 16 Liu W, Liu Z, Yao Z, et al. The prevalence and influencing factors of methicillin-resistant *Staphylococcus aureus* carriage in people in contact with livestock: a systematic review. *Am J Infect Control* 2015;43:469–75.
- 17 Voss A, Loeffen F, Bakker J, et al. Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerg Infect Dis* 2005;11:1965–6.
- 18 Bisdorff B, Scholthöfer JL, Claßen K, et al. MRSA-ST398 in livestock farmers and neighbouring residents in a rural area in Germany. *Epidemiol Infect* 2012;140:1800–8.
- 19 Schinasi L, Wing S, Augustino KL, et al. A case control study of environmental and occupational exposures associated with methicillin resistant *Staphylococcus aureus* nasal carriage in patients admitted to a rural tertiary care hospital in a high density swine region. *Environ Health* 2014;13:54.
- 20 Horgan M, Abbott Y, Lawlor PG, et al. A study of the prevalence of methicillin-resistant *Staphylococcus aureus* in pigs and in personnel involved in the pig industry in Ireland. *Vet J* 2011;190:255–9.
- 21 Angen Øystein, Feld L, Larsen J, et al. Transmission of methicillin-resistant *Staphylococcus aureus* to human volunteers visiting a swine farm. *Appl Environ Microbiol* 2017;83:e01489–17.
- 22 Bos MEH, Verstappen KM, van Cleef BAGL, et al. Transmission through air as a possible route of exposure for MRSA. *J Expo Sci Environ Epidemiol* 2016;26:263–9.
- 23 Cuny C, Nathaus R, Lauer F, et al. Nasal colonization of humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without exposure to pigs. *PLoS One* 2009;4:e6800.
- 24 Denis O, Suetens C, Hallin M, et al. Methicillin-resistant *Staphylococcus aureus* ST398 in swine farm personnel, Belgium. *Emerg Infect Dis* 2009;15:1098–101.
- 25 Garcia-Graells C, Antoine J, Larsen J, et al. Livestock veterinarians at high risk of acquiring methicillin-resistant *Staphylococcus aureus* ST398. *Epidemiol Infect* 2012;140:383–9.
- 26 Garcia-Graells C, van Cleef BAGL, Larsen J, et al. Dynamic of livestock-associated methicillin-resistant *Staphylococcus aureus* CC398 in pig farm households: a pilot study. *PLoS One* 2013;8:e65512.
- 27 Geenen PL, Graat EAM, Haenen A, et al. Prevalence of livestock-associated MRSA on Dutch broiler farms and in people living and/or working on these farms. *Epidemiol Infect* 2013;141:1099–108.
- 28 Graveland H, Wagenaar JA, Heesterbeek H, et al. Methicillin resistant *Staphylococcus aureus* ST398 in veal calf farming: human MRSA carriage related with animal antimicrobial usage and farm hygiene. *PLoS One* 2010;5:e10990.
- 29 Hatcher SM, Rhodes SM, Stewart JR, et al. The Prevalence of antibiotic-resistant *Staphylococcus aureus* nasal carriage among industrial hog operation workers, community residents, and children living in their households: North Carolina, USA. *Environ Health Perspect* 2017;125:560–9.
- 30 Huber H, Koller S, Giezendanner N, et al. Prevalence and characteristics of methicillin-resistant *Staphylococcus aureus* in humans in contact with farm animals, in livestock, and in food of animal origin, Switzerland, 2009. *Euro Surveill* 2010;15:19542.
- 31 Köck R, Harlizius J, Bressan N, et al. Prevalence and molecular characteristics of methicillin-resistant *Staphylococcus aureus* (MRSA) among pigs on German farms and import of livestock-related MRSA into hospitals. *Eur J Clin Microbiol Infect Dis* 2009;28:1375–82.
- 32 Lewis HC, Mølbak K, Reese C, et al. Pigs as source of methicillin-resistant *Staphylococcus aureus* CC398 infections in humans, Denmark. *Emerg Infect Dis* 2008;14:1383–9.
- 33 Moodley A, Nightingale EC, Stegger M, et al. High risk for nasal carriage of methicillin-resistant *Staphylococcus aureus* among Danish veterinary practitioners. *Scand J Work Environ Health* 2008;34:151–7.
- 34 Muters NT, Bieber CP, Hauck C, et al. Comparison of livestock-associated and health care-associated MRSA-genes, virulence, and resistance. *Diagn Microbiol Infect Dis* 2016;86:417–21.
- 35 Nadimpalli M, Stewart JR, Pierce E, et al. Livestock-associated, antibiotic-resistant *Staphylococcus aureus* nasal carriage and recent skin and soft tissue infection among industrial hog operation workers. *PLoS One* 2016;11:e0165713.
- 36 Neyra RC, Frisancho JA, Rinsky JL, et al. Multidrug-resistant and methicillin-resistant *Staphylococcus aureus* (MRSA) in hog slaughter and processing plant workers and their community in North Carolina (USA). *Environ Health Perspect* 2014;122:471–7.
- 37 Oppliger A, Moreillon P, Charrière N, et al. Antimicrobial resistance of *Staphylococcus aureus* strains acquired by pig farmers from pigs. *Appl Environ Microbiol* 2012;78:8010–4.
- 38 Pletinckx LJ, Verheghe M, Crombé F, et al. Evidence of possible methicillin-resistant *Staphylococcus aureus* ST398 spread between pigs and other animals and people residing on the same farm. *Prev Vet Med* 2013;109:293–303.
- 39 Richter A, Sting R, Popp C, et al. Prevalence of types of methicillin-resistant *Staphylococcus aureus* in turkey flocks and personnel attending the animals. *Epidemiol Infect* 2012;140:2223–32.
- 40 Rinsky JL, Nadimpalli M, Wing S, et al. Livestock-associated methicillin and multidrug resistant *Staphylococcus aureus* is present among industrial, not antibiotic-free livestock operation workers in North Carolina. *PLoS One* 2013;8:e67641.
- 41 Sahibzada S, Hernández-Jover M, Jordan D, et al. Emergence of highly prevalent CA-MRSA ST93 as an occupational risk in people working on a pig farm in Australia. *PLoS One* 2018;13:e0195510.
- 42 Schmitthausen RM, Schulze-Geisthoevel SV, Stemmer F, et al. Analysis of transmission of MRSA and ESBL-E among pigs and farm personnel. *PLoS One* 2015;10:e0138173.
- 43 Van Cleef BAGL, Broens EM, Voss A, et al. High prevalence of nasal MRSA carriage in slaughterhouse workers in contact with live pigs in the Netherlands. *Epidemiol Infect* 2010;138:756–63.
- 44 van Cleef BAGL, van Benthem BHB, Verkade EJM, et al. Dynamics of methicillin-resistant *Staphylococcus aureus* and methicillin-susceptible *Staphylococcus aureus* carriage in pig farmers: a prospective cohort study. *Clin Microbiol Infect* 2014;20:0764–71.
- 45 van Cleef BAGL, van Benthem BHB, Verkade EJM, et al. Livestock-associated MRSA in household members of pig farmers: transmission and dynamics of carriage, a prospective cohort study. *PLoS One* 2015;10:e0127190.
- 46 VAN DEN Broek IVF, VAN Cleef BAGL, Haenen A, et al. Methicillin-resistant *Staphylococcus aureus* in people living and working in pig farms. *Epidemiol Infect* 2009;137:700–8.
- 47 van Duijken E, Hengeveld P, Zomer TP, et al. Transmission of MRSA between humans and animals on duck and turkey farms. *J Antimicrob Chemother* 2016;71:58–62.
- 48 van Loo I, Huijsdens X, Tiemersma E, et al. Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans. *Emerg Infect Dis* 2007;13:1834–9.
- 49 Vandendriessche S, Vanderhaeghen W, Soares FV, et al. Prevalence, risk factors and genetic diversity of methicillin-resistant *Staphylococcus aureus* carried by humans and animals across livestock production sectors. *J Antimicrob Chemother* 2013;68:1510–6.
- 50 Verkade E, Kluytmans-van den Bergh M, van Benthem B, et al. Transmission of methicillin-resistant *Staphylococcus aureus* CC398 from livestock veterinarians to their household members. *PLoS One* 2014;9:e100823.
- 51 Walter J, Espelage W, Adlhoef C, et al. Persistence of nasal colonisation with methicillin resistant *Staphylococcus aureus* CC398 among participants of veterinary conferences and occurrence among their household members: a prospective cohort study, Germany 2008–2014. *Vet Microbiol* 2017;200:13–18.
- 52 Walter J, Espelage W, Cuny C, et al. Veterinarians visiting swine farms are at high risk for colonization with livestock-associated methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 2016;62:126.1–8.

- 53 Wulf MWH, Tiemersma E, Kluytmans J, *et al.* MRSA carriage in healthcare personnel in contact with farm animals. *J Hosp Infect* 2008;70:186–90.
- 54 Crombé F, Argudin MA, Vanderhaeghen W, *et al.* Transmission dynamics of methicillin-resistant *Staphylococcus aureus* in pigs. *Front Microbiol* 2013;4:57.
- 55 Porphyre T, Giotis ES, Lloyd DH, *et al.* A metapopulation model to assess the capacity of spread of methicillin-resistant *Staphylococcus aureus* ST398 in humans. *PLoS One* 2012;7:e47504.
- 56 Feingold BJ, Silbergeld EK, Curriero FC, *et al.* Livestock density as risk factor for livestock-associated methicillin-resistant *Staphylococcus aureus*, the Netherlands. *Emerg Infect Dis* 2012;18:1841–9.
- 57 Topp E. Agriculture and agri-food Canada's research program on antimicrobial resistance. *Can Commun Dis Rep* 2017;43:224–7.
- 58 Espinosa-Gongora C, Broens EM, Moodley A, *et al.* Transmission of MRSA CC398 strains between pig farms related by trade of animals. *Vet Rec* 2012;170:564.
- 59 Grøntvedt CA, Elstrøm P, Stegger M, *et al.* Methicillin-resistant *Staphylococcus aureus* CC398 in humans and pigs in Norway: a "one health" perspective on introduction and transmission. *Clin Infect Dis* 2016;63:1431–8.
- 60 Smith TC. Livestock-associated *Staphylococcus aureus*: the United States experience. *PLoS Pathog* 2015;11:e1004564.