and cancer incidence. Personal air styrene measurements and biological markers of styrene exposure from the 1960s up to the present day will be used to update the exposure assessment. Linear mixed-effects models will be applied to develop a quantitative, historical, industry-specific job-exposure matrix. Predictors available in the individual cohorts include country, occupation, employment year, product, process, and task. Some cohorts will be pooled together prior to analysis, others analyzed separately following a common protocol that will focus on different exposure metrics (cumulative, duration, mean, highest attained, peaks) and exposure time windows. The aggregated data will be synthesized by a meta-analysis.

Results A total of 96,000 workers employed between 1947–2007 in 762 companies in Finland, Italy, UK, US, and Denmark, and over 2.8 million person years will be included, and 40,000 air samples and 13,000 urinary samples are identified.

Conclusion Pooling and meta-analysis of existing cohorts are powerful tools in the search for more definite answers to the carcinogenicity of this important chemical.

P-323

THE EFFECT OF THE ONSET OF A DISEASE ON EXIT FROM PAID EMPLOYMENT AMONG WORKERS IN THE NETHERLANDS: A LONGITUDINAL REGISTER-BASED STUDY WITH 11 YEARS FOLLOW-UP

¹Roos Hijdra, Suzan Robroek, Lex Burdorf, Merel Schuring. ¹Erasmus Medical Center, Netherlands

10.1136/OEM-2021-EPI.270

Introduction With increasing retirement age, employed persons are more likely to suffer from chronic diseases, such as cardiovascular diseases, diabetes, inflammatory diseases, respiratory diseases, and psychological disorders, at later stage during their working career.

Objectives This study investigates the influence of the onset of a disease on the likelihood of exit from paid employment through different pathways (unemployment, disability benefits, economic inactivity, early retirement).

Methods For this longitudinal study with 11 years of follow-up (2009–2019), 4,276,770 persons were selected, who were employed between 2009 and 2010, and used no medication for the selected diseases in 2009. Register data from Statistics Netherlands on medication use, employment status, and socio-demographic factors were used. Exit from paid employment pathways were defined as: unemployment, disability benefits, economic inactivity, and early retirement. The following six disease categories were identified based on medication: cardiovascular diseases, diabetes mellitus, respiratory illness, psychological disorders, inflammatory disorders and psychotic disorders. Descriptive statistics and Cox Proportional Hazards analyses with competing risks were performed.

Results The onset of any disease increased the likelihood of exit from paid employment, with strongest effect observed for psychological disorders (Hazard Ratio [HR]=1.37, 95% confidence interval [CI] 1.34–1.40) and psychotic disorders (HR=1.49, 95% CI 1.41–1.58). Of all pathways out of paid employment, onset of a disease had the strongest effect on disability benefits, with HRs ranging from 1.75 (inflammatory diseases) to 9.30 (psychotic disorders).

Conclusion For the working age population, the onset of common disorders, especially psychological and psychotic disorders, is a risk for maintaining paid employment. Interventions

are needed to prevent persons with these diseases from involuntary loss of paid employment.

P-328

AGE DIFFERENCES IN WORK-DISABILITY DURATION ACROSS CANADA: EXAMINING VARIATIONS BY FOLLOW-UP TIME AND CONTEXT

¹Jonathan Fan, Robert Macpherson, Peter Smith, Anne M Harris, Monique Gignac, Chris McLeod. ¹University of British Columbia, Canada

10.1136/OEM-2021-EPI.271

Introduction Research has examined age-related patterns in return to work and wage-replacement duration following a workplace injury. The various clinical, functional or physiological factors studied do not fully account for age differences in wage-replacement duration. One contextual factor that has been largely overlooked in research studies is the potential impact of the phase of recovery.

Objectives This study aimed to understand age differences in wage-replacement duration by focusing on variations in the relationship across different periods of follow-up time.

Methods We used administrative claims data provided by six workers' compensation systems in Canada, focusing on time-loss claims for workers aged 15–80 years with a work-related injury/illness during the 2011 to 2015 period. Survival analysis examined age-related differences in the hazard of transitioning off (versus remaining on) disability benefits, allowing for relaxed proportionality constraints on the hazard rates over time. Differences were examined on the absolute (hazard difference) and relative (hazard ratios [HR]) scales.

Results Older age groups had a lower likelihood of transitioning off wage-replacement benefits compared to younger age groups in the overall models (e.g., 55–64 vs. 15–24 years: HR 0.62). However, absolute and relative differences in age-specific hazard rates varied as a function of follow-up time. The greatest age-related differences were observed at earlier event times and were attenuated towards a null difference across later follow-up times.

Conclusion Our study provides insight into the workplace injury/illness claim and recovery processes and suggests that older age is not always strongly associated with worse disability duration outcomes at longer disability durations. The use of data from multiple jurisdictions lends external validity to our findings and demonstrates the utility of using cross-jurisdictional data extracts. Future work should examine the social and contextual determinants that operate during various recovery phases, and how these factors interact with age.

P-330

AGE DIFFERENCES IN RETURN-TO-WORK FOLLOWING INJURY: UNDERSTANDING THE ROLE OF AGE ACROSS LONGITUDINAL FOLLOW-UP

¹Jonathan Fan, Monique Gignac, Anne M Harris, Peter Smith. ¹University of British Columbia, Canada

10.1136/OEM-2021-EPI.272

Introduction Older age tends to be associated with longer time to return-to-work (RTW) following a workplace injury and multiple recurrences of work absence following an initial RTW attempt. However, few studies have examined the underlying factors that are responsible for these differences.