disinfecting chemicals in the workplace ($p = 0.01$; OR = 17.306) and the absence of occupational reclassification ($p = 0.01$; OR = 21.567).

Conclusion This study has revealed skin-related QoL is significantly more impaired in women. Several studies have reported a similar result. This could be explained by the fact that traditionally, in Tunisia, women have been predominantly responsible for household chores, and therefore more exposed to substances that can trigger dermatological conditions. In addition, the aesthetic consequences of skin pathology do not have the same impact on men and women.

P-363 SEX DIFFERENCES IN MANAGING ACCIDENTAL BLOOD EXPOSURE IN A TUNISIAN PUBLIC HOSPITAL

¹Amira Omrane, Harrathi Chayma, Meriem Ben Hassine, Lamia Bouzgarrou, Aouatef Mahfoudh, Taoufik Khalfallah. ¹Faculty of Medicine of Monastir, Tunisia

10.1136/OEM-2021-EPI.290

Accidental blood exposure (ABE) is major problem of occupational safety among health care workers. This work aimed to assess the gender differences among health care workers of a public hospital in the central region of Tunisia.

Methods An exhaustive retrospective study was conducted over a period of fourteen years, concerning ABE reported in the Department of Occupational Medicine of a public hospital in the central region of Tunisia (Mahdia). The data collected were related to socio-professional characteristics of victims, circumstances of the accident and immediate and follow-up care. Gender groups were compared.

Results A total of 650 ABE were reported during the study period. Victims were predominately female (sex ratio=0.47), mean aged 32.1 ± 9.4 years and having a mean job tenure of 5.5 ± 7.7 years. Paramedics represented 28.3%. AEB were caused by a needle stick in 82.6% of cases. Needle recapping was the direct cause in 10.6% of cases. Only 47.1% of caregivers were wearing gloves at the time of ABE. Women were more exposed to ABE with a higher risk of contamination, but with no statistically significant difference ($p=0.33$). Immediate care was in accordance with universal recommendations in 34.1% of cases and less adequate among women ($p=0.05$). The initial serology was prescribed in 82.6% of cases, that of the 3rd month in 5% of cases and that of the 6th month in 1.2% of cases. Women were less adherent to follow-up with a statistically significant difference at six months ($p=0.016$). Hepatitis B vaccination schedule was correctly followed in 79% of cases. Women were more adherent than men ($p=0.029$). Logistic regression analysis did not reveal gender as a predictor of vaccination status ($p=0.211$).

Conclusion The current study has revealed a high prevalence of ABE mainly among female health care workers with unsatisfactory management, including inadequate immediate care and incomplete post-accident follow-up.

P-364 INTERACTIONS AMONG SHIFT WORK DOMAINS AND EMPTY CALORIE FOOD/BEVERAGE CONSUMPTION

¹Ting-Ti Lin, Shannon Zenk, Pamela Martyn-Nemeth, Lisa Tussing-Humphreys, Kathleen M Rospenda, Chang Park, Mary C Kapella. ¹National Defense Medical Center, Taiwan

10.1136/OEM-2021-EPI.291

Introduction Shift work has been linked to increased consumption of empty calorie food/beverages. However, the majority of studies investigating associations between shift work and empty calorie food/beverage consumption has been focusing on the impacts of shift timing. Little is known about how other domains of shift work contribute to empty calorie food/beverage consumption.

Objectives The purpose of this study was to examine associations between shift work and empty calorie food/beverage consumption, focusing on other domains of shift work and their interactions.

Methods This was a 14-day intensive longitudinal study with ecological momentary assessment. A convenience sample of 80 Taiwanese hospital registered nurses were recruited. During the study period, employing a 21-item food checklist, participants were prompted four times daily to report their empty calorie food/beverage consumption on a smartphone. Three domains of shift work: shift timing, intensity, and speed were derived from the registry-based work schedules. Three-level mixed-effects regression models were used to test hypotheses.

Results A total of 2,444 momentary observations from 77 participants were included in the final analysis. Findings suggested that high night shift intensity was associated with an increased likelihood of sugar-sweetened beverage intake (AOR=1.64, 95% CI [1.01, 2.68]). Shift speed moderated associations between sugar-sweetened beverage consumption and work shift intensity or shift timing. However, associations between night shift intensity and empty calorie food/beverage consumption did not vary by shift speed.

Conclusions This study suggested how shift assignments might contribute to workers' empty calorie food/beverage consumption. Therefore, it would be beneficial to rotating shift workers' eating behaviors and overall health if the identified hazardous shift schedule can be avoided.

P-365 PROMOTING SENTINEL SURVEILLANCE PROGRAMS FOR ANTIMICROBIAL RESISTANCE IN CANADIAN SALMON AQUACULTURE, A POSSIBLE AND UNDERSTATED OCCUPATIONAL HEALTH HAZARD

¹Cory Ochs, Barb Neis, Kapil Tahlan, Atanu Sarkar. ¹Memorial University of Newfoundland and Labrador, Canada

10.1136/OEM-2021-EPI.292

Introduction Canadian salmon aquaculture is a high-risk industry with injury rates that surpass provincial averages in Atlantic Canada, yet few publications address occupational health hazards. Antimicrobial resistance (AMR) emergence is a growing public health concern, and the marine aquatic environment with its rich microbiota is particularly vulnerable to selection of AMR. Antibiotic use in the industry and other anthropogenic activities that result in the deposit of pollutants contaminated with antibiotics into the marine environment can together amplify selective pressure propagating AMR. Similar to terrestrial animal production facilities, there is concern for the development of hotspots for occupational exposures to AMR among aquaculture workers. As the fastest growing food production network globally, the aquaculture industry has been appealed by the Food and Agriculture Organization, among others, to standardize monitoring and to generate an evidence base to better understand the aetiology of AMR emergence in aquaculture settings.