significantly associated with RCD surgery after more than 10 years of work (Ex. HR for 11–20 years vs. 0 years with static strength score $\geq 4 = 2.06$, 95%CI=1.39–3.04).

Conclusion Numerous occupational physical exposures were associated with incident RCD surgery. Associations were strongest in workers with more than a decade of high exposure.

0-240

ASSOCIATIONS BETWEEN FEELING COLD AT WORK AND WORK PERFORMANCE IN A COLD-EXPOSED WORKING POPULATION FROM THE TROMSØ 6 STUDY

¹Morten Skandfer, Erlend Farbu Hoftun, Tormod Brenn, Anje Christina Höper. ¹UIT The Arctic University of Norway, Norway

10.1136/OEM-2021-EPI.87

Objective Cold exposure is associated with an increased prevalence of musculoskeletal pain. We found earlier that employees spending $\geq 25\%$ of their working time in cold environments had higher odds of chronic musculoskeletal pain. There was a consistent tendency of higher odds for increasing frequency of feeling cold. Cold exposure can also interfere with work performance. The aim of this study was therefore to investigate if the frequency of cold experience was associated with impaired work performance.

Methods We used data from the sixth survey of the Tromsø study (2007–2008). Participants aged 30–67 years who reported to work in a cold environment $\geq 25\%$ of the time, were not retired, not receiving full-time disability benefits and without missing values were included, leading to 793 participants. Feeling cold was categorized into never, sometimes and often feeling cold. Work performance variables comprised of binary variables of impaired control of movement, heavy physical work and long-lasting physical work, finger dexterity and –sensitivity. Associations between feeling cold at work and self-reported work performance were examined with Poisson-regression, adjusted for age, sex, smoking and body mass index.

Results Both prevalence of impaired work performance and associations between frequency of feeling cold and impaired work performance were consistently lower for those never feeling cold and higher for those feeling cold often, compared to those feeling cold sometimes. In the fully adjusted model, the strongest associations were found for impaired long-lasting work performance with prevalence ratio (PR) 0,35 (95%CI 0,20–0,62) for never feeling cold and PR 1,81 (95%CI 1,35–2,42) for feeling cold often. For impaired heavy work PRs were 0,53 (95%CI 0,31–0,90) and 2,13 (95%CI 1,50–3,04), respectively.

Conclusion In this cross-sectional study on cold-exposed workers, cold experience frequency was associated with work performance with the tendency of increased work impairment with increasing frequency of feeling cold.

0-433

WHICH QUEBEC INDUSTRIES AND OCCUPATIONS ARE AT RISK OF WORK-RELATED MUSCULOSKELETAL DISORDERS? A COMPARISON OF ANALYSES OF 2010–2012 WORKERS' COMPENSATION AND 2014–2015 HEALTH SURVEY DATA

¹Susan Stock, Nektaria Nicolakakis, France Tissot. ¹INSPQ; University of Montreal, Canada

10.1136/OEM-2021-EPI.88

Introduction Non traumatic work-related musculoskeletal disorders (WMSD) represent an enormous burden of preventable illness. Two strategies and data sources to document this burden and identify workers at highest risk were compared.

Objectives To identify gender-stratified worker groups at high risk of non-traumatic WMSD by industry and type of occupation and compare WC to health survey results.

Methods Using 2014–2015 Quebec Health Survey (QPHS) data on 24,300 workers, measuring self-reported WMSD and industry groupings stratified by occupation (manual/mixed/non-manual), WMSD risk for each industry-occupation group was estimated using gender-stratified adjusted regression analyses and estimation methods. Using Quebec 2010–2012 workers' compensation (WC) data, gender-stratified WMSD incidence rates per 1,000 full-time equivalent employees (% FTEE) were calculated for 174 industry-type-of-occupation groups. WMSD risk was ranked according to Prevention Index scores.

Results In both studies, women in manual occupations had the highest WMSD risk compared to male counterparts (WC: 39%vs27% FTEE; QPHS: 36%vs25%); manual male and female workers in administrative/support/cleaning/garbage services were identified at high risk; as well as women in accommodation/restaurant and men in specialised construction trades, civil engineering, and metal manufacturing. Compensation data identified another 9 high-risk groups for men, and 11 for women including 3 health sector groups that ranked in the top 5 for women. Conversely, the QPHS identified another 13 high risk groups in men including several construction and manufacturing sectors and 5 in women.

Discussion Differences between the 2 studies' results are likely due to methodologic differences, including under-reporting in compensation data and the survey's low power to identify some industries stratified by gender and occupation. Results of the two studies are complementary and each adds to our understanding of which groups are at WMSD risk to target for prevention. Research is needed to compare different survey and compensation data analytic strategies to improve capacity to identify workers at high WMSD risk.

0-465

PAIN IN HEALTHCARE WORKERS: A PERSPECTIVE OF MULTIDISCIPLINARY APPROACH

¹Simone Fargetti, Telma de Cassia dos Santos Nery, Salvador Celso Callia, Maria Thereza Toledo Penteado, Monize Mendonça Cruz, Moacyr Vergara de Godoy Moreira. ¹Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, Brazil

10.1136/OEM-2021-EPI.89

Introduction Health professionals deal with numerous hazards during their occupational life from a physical and mental point of view: precarious working conditions, irregular working hours, emotional pressure, physically demanding jobs, involving weight lifting and manual tasks, which may cause pain and musculoskeletal disorders. Our occupational health service assists more than 20,000 health workers in a large hospital complex in Latin America. It consists of a multidisciplinary team involving physicians, psychologists, engineers, occupational therapists, physiotherapists, as well as an emergency care service where workers may seek immediate medical attention.

Objective To describe the most frequent complaints related to musculoskeletal pain in emergency medical care at our

occupational medicine service and the multidisciplinary approach for referring and treating these workers.

Methods This descriptive and retrospective study comprised data from the occupational emergency care service from 2016 to 2020. All complaints related to International Classification of Diseases (ICD10) codes M and S were analyzed.

Results 13,312 consultations were carried out in our service from 2016 to 2020 due to ICDs M and S; 73.1% workers were female, which coincides with genre distribution in our service. The leading cause of musculoskeletal pain was low back pain (M54.5), whereas finger injuries (S61.0) and ankle sprains (S93.0) were the most important complaints related to trauma. Visits related to musculoskeletal complaints were less frequent in 2020 compared to other years due to Covid-19 pandemics, since this service was responsible for evaluating and testing workers with respiratory symptoms. Workers with musculoskeletal complaints were referred directly from emergency care or occupational physician to multidisciplinary team: orthopedic surgeon, acupuncture, physiotherapy or work adjustment with occupational team.

Conclusion Occupational emergency care data is an important indicator of injuries and pain in healthcare professionals. Its integration with a multidisciplinary team is essential to prevent further musculoskeletal illnesses in hospital workforce.

Musculoskeletal-2

O-20 PREVALENCE, PREDICTORS AND WAGE REPLACEMENT DURATION ASSOCIATED WITH DIAGNOSTIC IMAGING IN AUSTRALIAN WORKERS WITH ACCEPTED CLAIMS FOR LOW BACK PAIN: A RETROSPECTIVE COHORT STUDY

¹Michael Di Donato, Ross Iles, Rachelle Buchbinder, Ting Xia, Alex Collie. ¹Monash University, Australia

10.1136/OEM-2021-EPI.90

Introduction Diagnostic imaging is not recommended for low back pain (LBP) in the absence of clinical evidence to suggest a serious pathology is the cause of pain. Workers may access funding for wage replacement and healthcare, including diagnostic imaging, from workers' compensation if they cannot work due to LBP.

Objectives This study sought to determine in Australian workers with accepted workers' compensation claims for LBP (1) the prevalence of diagnostic imaging of the spine and factors associated with its use, and (2) the association between spinal diagnostic imaging events and wage replacement duration.

Methods Workers with accepted workers' compensation claims for LBP longer than two weeks were grouped by whether workers' compensation funded no, single, or multiple diagnostic spinal imaging in the two years since reported low back pain onset. Ordinal logistic regression was used to define the demographic, occupational and social factors associated with each group. Time-to-event analysis was used to determine the association between spinal imaging and wage replacement duration.

Results In the sample of 30,530 workers, 9,267 (30.4%) received single spinal imaging and 6,202 (20.3%) received multiple spinal imaging. Male workers and workers from the

state of Victoria had significantly higher odds of multiple imaging. Socioeconomically advantaged workers and workers from remote Australia had significantly lower odds of multiple imaging. Magnetic Resonance Imaging was the most common imaging modality. Workers with single spinal imaging (median duration 17.0 weeks; HR 2.0, 95%CI 1.9, 2.1) and multiple spinal imaging (median duration 49.0 weeks; HR 4.0, 95%CI 3.9, 4.1) had significantly longer wage replacement duration than those with no imaging (median duration 6.1 weeks).

Conclusions Over half of Australian workers with an accepted workers' compensation claim for LBP longer than two weeks received diagnostic spinal imaging. Receipt of diagnostic imaging, particularly multiple imaging, was associated with longer wage replacement duration.

O-202 WORKERS ON PROLONGED WORK DISABILITY FOR MUSCULOSKELETAL DISORDERS DO NOT WORRY FOR NOTHING

¹Marie-France Coutu, Marie-José Durand, Fergal O'Hagan, Patrick Gosselin, Iuliana Nastasia, Djamel Berbiche, Marie-Elise Labrecque, Sara Pettigrew. ¹Université de Sherbrooke, Canada

10.1136/OEM-2021-EPI.91

Introduction Workers worry during prolonged work disability, but do their worries relate to their actual work disability situation?

Objective The aim of this study was to assess worries and their maintaining factors, while considering the margin of maneuver/leeway at work and their impact on return to work (RTW).

Methods We conducted a cohort study with a convenience sample of 79 (39 men and 40 women) workers having persistent (≥ 3 months) work-related musculoskeletal disorders causing absence from their regular work. Following Dugas' theory validated self-administered questionnaires (ex.: intolerance of uncertainty, utility of worrying) were completed at the beginning and the end of the work rehabilitation program. Also, the questionnaire on type of worries (QTW) assessed specific types of worries and their relationship to work. Trained occupational therapists, (n = 16) evaluated the margin of maneuver of all workers. Multivariate analyses were performed on RTW predicted by workers' indicators and occupational therapists' margin of maneuver.

Results

Twenty-one workers did not RTW The model predicted 54% of the variance in N-RTW (p .0001). Significant factors explaining N-RTW were: lack of a margin of maneuver (OR = 8.5; p = .008); high intolerance for uncertainties (OR = 1.12; p = .01), perceived utility of worrying (OR = 1.11; p .001), and for the QTW scores, a high mean intensity of worries (OR = 2; p = .004) emerging from actual situations (OR = 17.15; p = .02) occurring at work (OR = 8.5). A posthoc analysis (pseudo R² = .33; p =) shows that a lack of a margin of maneuver is associated with QTW scores of worries emerging from uncertainties at work.

Conclusion Workers not returning to work worry about actual situations at work, but this is also associated with low margin of maneuver, assessed by occupational therapists. Thus, RTW interventions should focus on the work environment.