(AOR=2.37, 95% CI= 1.26–2.93) each significantly related to increased odds of verbal violence. Short duration of employment (<five years) was the most strongly associated factor for sexual harassment (AOR= 2.82, 95% CI=1.7–2.94). Working the night shift was significantly associated with encountering verbal violence (AOR= 2.79, 95% CI= 1.69–3.06).

Conclusion The findings of this study highlighted occupational violence and the need for an awareness of violence against female workers in industrial plants. The high prevalence of violence suggests that management in industrial workplaces should invest time and resources in addressing violence. Providing reporting procedure and training programs are recommended in dealing with workplace violence risk.

O-158 LONG-TERM PROGNOSIS FOR HEALTH CARE UTILIZATION AMONG PATIENTS WITH WORK-RELATED MENTAL DISORDERS

¹Morten Vejs Willert, Marianne Kyndi, Vita Ligaya Dalgaard, David Høyrup Christiansen, Johan Hviid Andersen. ¹Aarhus University Hospital, Denmark

10.1136/OEM-2021-EPI.114

Objective The long-term prognosis for employees with work-related stress, depression or PTSD is unclear. We aim to investigate if these conditions represent a temporary setback or a decline towards long-term poor health.

Methods From the Danish Occupational Medicine Cohort we included patients referred 2000–2012 with mental disorders (N=17.900) to Departments of Occupational Medicine across Denmark. From nation-wide registries we obtained data on general health care service utilization (GP contacts) and specialized mental health care (incident private practice psychiatrist/psychologist care and psychiatric hospital care) for five years after/before the index-date of referral. Results are presented as utilization 5 years before referral (index-year -5), the year on either side of referral (index-year -1 and +1) and five years after referral (index-year +5).

Results General health care utilization increased from 7 median GP contacts in index-year -5 rising sharply to 12 in index-year -1 and 10 in index-year +1, then declining gradually to 8 in index-year +5. Incident private practice psychiatrist/psychologist care was 4.5% among patients in index-year -5 rising to 13.6% in index-year -1, 19.8% in index-year +1, declining to 7.4% in index-year +5. Incident psychiatric hospital care was 1.4% in index-year -5 rising to 5.6% in index-year -1, 8.6% in index-year +1, then declining to 4.7% in index-year +5.

Conclusions General health care utilization was elevated in the years surrounding referral, then declined to levels comparable to five years prior to referral. From a relatively low incidence, mental health care utilization increased 4–6 fold in the years around referral, then declined to levels 2–3 fold higher than five years before referral. This could indicate that work-related mental disorders represent a temporary setback for most patients, but for some becomes a decline towards long-term poor mental health. Further analyses will seek to identify at-risk groups from demographic, occupational and health-related prognostic factors.

O-189 MENTAL HEALTH IN CHILDHOOD AND ADOLESCENCE MATTERS: LINKING MENTAL HEALTH TRAJECTORIES WITH EMPLOYMENT OUTCOMES IN YOUNG ADULTHOOD

¹Marlene Amalie Magerl, Karin Veldman, Ute Bültmann, Iris Arends. ¹Faculty of Medical Sciences, University Medical Center Groningen and University of Groningen, Netherlands

10.1136/OEM-2021-EPI.115

Introduction Mental health problems cause a considerable burden of disease in adolescents, with potential negative consequences on employment later in life. Earlier, five trajectories of internalising and externalising problems (high-stable, moderate-high, decreasing, moderate-low, low-stable) among adolescents from 11 to 19 years were identified and a relationship with employment status at age 19 was found. However, at age 19, many participants were still in education.

Objectives This study aims to examine associations of trajectory membership with having paid work and type of employment contract in young adults at age 26.

Methods Fifteen-year follow-up data of the longitudinal TRacking Adolescents' Individual Lives Survey (TRAILS) cohort study (N=1711) were used. Logistic regression analyses were applied to examine associations of trajectory membership of internalising and externalising problems with having paid work and type of employment contract at age 26. The analyses were stepwise adjusted for sex, intelligence, parental education, family composition, physical health, negative life events and mental health comorbidity, all measured at age 11.

Results For internalising problems, the high-stable (adjusted OR: 2.19; 95% CI: 1.17–4.11) and moderate-high (adjusted OR: 1.75; 95% CI: 1.14–2.69) trajectories were strongly associated with not having paid work, compared to the low-stable trajectory. No associations were found between the trajectories of externalising problems and having paid work. For both internalising and externalising trajectories, no associations were found between trajectory membership and having a permanent or temporary employment contract.

Conclusion This study expands current knowledge about the impact of internalising problems on employment of young adults. To prevent the effects of mental health problems on work, a life course perspective is needed taking into account the development of mental health problems in childhood and adolescence. In the transition from school to work, integrating youth- and occupational healthcare may be key to prevent negative employment outcomes due to internalising problems.

O-211 OCCUPATIONAL DAYTIME LIGHT EXPOSURE AND RISK OF INITIATING ANTIDEPRESSANT TREATMENT AMONG MEN AND WOMEN: A NATIONAL FOLLOW-UP STUDY

¹Anne Vested, Vivi Schlünssen, Lex Burdorf, Esben Meulengracht Flachs, Stine Daugaard, Jens Peter Ellekilde Bonde, Susan Peters, Anne Helene Garde, Zara Ann Stokholm, Morten Vejs Willert, Jesper Medom Vestergaard, Henrik Albert Kolstad. ¹Aarhus University, Denmark

10.1136/OEM-2021-EPI.116

Introduction Higher winter depression prevalence at temperate latitudes has been attributed to low daylight levels. Daytime light exposure varies across jobs.

Objectives In a nationwide study, we examined occupational daytime light exposure levels and risk of initiating antidepressant treatment among men and women.

Methods We conducted a register-based follow-up study of all gainfully employed 526 839 men and 506 092 women in Denmark (56° N) born 1980 or later. We estimated annual mean occupational daytime (6:00–17:59 h) white light exposure levels (lux) for each year of employment 2000–2015 with a quantitative job exposure matrix (JEM). We followed participants from 2001 or first year of employment (if later) until first antidepressant treatment $\leq$ 2016 as recorded in a national prescription database. We calculated incidence rate ratios adjusted for register information on age, calendar year, family history of antidepressant treatment, and educational level. Smoking was assessed with a lifestyle JEM.

Results During follow up, the population accumulated 6 804 361 person-years and 30 816 men (5.8%) and 52 817 women (10.4%) initiated antidepressant treatment corresponding to 8 and 16 cases per 1000 person-years, respectively. Two percent of women and eight percent of men were exposed to an annual average above 500 lux. Among women, we observed reduced risk of initiating antidepressant treatment for those exposed the previous year to 500–999 lux (incidence rate ratio [IRR] = 0.92, 585 cases, 95% CI 0.85–1.00) and $\geq$ 1000 lux (IRR = 0.90, 294 cases, 95% CI 0.80–1.01) when compared with $<$ 250 lux. No association was apparent for men.

Conclusion Our results indicate a preventive effect of daytime light exposure on depression and other mental disorders treated with antidepressants for the small fraction of workers exposed to an annual mean above 500 lux at work, but only among women. Findings need replication in other countries.

O-359

MENTAL HEALTH IN A SAMPLE OF 72,452 WORKERS FROM 8 COUNTRIES OF LATIN AMERICA: RESULTS OF HEALTH AND WORKING CONDITIONS SURVEYS

¹Paula Santia, Michael Silva-Peñaherrera, Fernando G Benavides. ¹Universitat Pompeu Fabra, Spain

10.1136/OEM-2021-EPI.117

Introduction Work is considered a key determinant of mental health. However, evidence on the impact of work in mental health in Latin America is scarce.

Objectives To analyse the relationship between mental health status and work in Latin America through three indicators: labour relationship (employed/self-employed or employer), employment condition (formal/informal), and occupational category (non manual skilled/non manual non-skilled/manual skilled/manual non-skilled).

Methods Cross sectional study based on health and working condition surveys from 8 countries of Latin America. Poor mental health was defined as a score of ≥ 3 on the GHQ12; ≥ 10 on the PHQ9; or ≤ 13 on the WHO5 scales. Informal workers were defined as those unregistered or lacking a contract. Occupational categories were created following the international standard classification of occupation. Prevalence ratios and their IC95% were calculated stratified by sex and adjusted by age, educational level and marital status.

Results Final sample consisted of 72,452 workers, from Argentina (n=8966), Brazil (n=52832), Costa Rica (n=1503), El Salvador (n=1507), Guatemala (n=1510), Honduras (n=1507), Nicaragua (n=1500) and Panamá (n=1505). Poor

mental health ranged from 4.5% in Brazilian men to 25.9% in Panamanian women. Manual non-skilled men showed higher prevalence of poor mental health than non-manual skilled men in Argentina (RPa 1.43 IC95% 1.04 - 1.95) and Central America (RPa 1.94 IC95% 1.12 - 3.34). In Brazil, men employers showed higher prevalence of poor mental health (RPa 1.25 IC95% 1.02 - 1.53) and informal workers showed higher prevalence of poor mental health irrespective of sex (Men RPa 1.16 IC95% 1.02 - 1.33; Women RPa 1.30 IC95% 1.05 - 1.60).

Conclusion Prevalence of poor mental health varies significantly between occupational groups and countries. Special attention should be pay to informal workers, workers in elementary occupations and employers.

Radiation

O-138

CANCER MORTALITY THROUGH 2016 AMONG URANIUM MINERS ON THE COLORADO PLATEAU: INTERACTIONS OF RADON WITH SMOKING AND TIME SINCE EXPOSURE

¹Mary Schubauer-Berigan, Stephen Bertke, Kaitlin Kelly-Reif, Robert Daniels. ¹International Agency for Research on Cancer, France

10.1136/OEM-2021-EPI.118

Objective The long-term risks of lung cancer from radon progeny exposure are known largely from studies of uranium miners, including a cohort of 4137 male miners from the Colorado Plateau (CP) in the southwestern US. Our objective was to estimate the excess absolute risk (EAR) and excess relative risk (ERR) of lung and other cancer from radon exposure among CP uranium miners, evaluating the persistence of risk with 26 years of extended follow-up, and the form of interaction between radon and cigarette pack-years (PY).

Methods We followed the CP cohort for mortality through 2016, re-evaluated its radon exposure against original records, and imputed a smoking end date for cohort members. We evaluated EAR based on standardized rate ratios of lung cancer, adjusted for age, calendar year, and race. We modeled ERR using Cox proportional hazards regression of full risk sets (adjusted for age, race, birth year, exposure rate and time since last exposure). We also modeled variation in risk per unit exposure by attained age (AA) and time-since-exposure (TSE) and evaluated joint effects of radon and PY.

Results We identified 695 cases of lung cancer, including 146 among never- and light smokers. Lung cancer EAR per unit radon exposure increased substantially with smoking PY and throughout decades of follow-up. Lung cancer ERR decreased with AA and TSE, although to a lesser extent than observed elsewhere. The form of interaction between radon and PY was sub-multiplicative but greater-than-additive, appearing closer to multiplicative at lower radon exposures. Pancreas was the only other site showing a significantly positive ERR per unit exposure, although much lower than for lung cancer.

Conclusion Excess risk of lung cancer persists throughout the lifespan among this cohort of uranium miners. Information about radon-smoking interactions, particularly at low exposure levels, is of interest for extrapolations to the general population.