O-461

OCCUPATIONAL SOLAR ULTRAVIOLET RADIATION AND BREAST CANCER RISK IN CANADA¹Sareena McDonald, Cheryl Peters, Dylan O'Sullivan, Will King, Darren Brenner. ¹Queen's University, Canada

10.1136/OEM-2021-EPI.15

Introduction Solar ultraviolet radiation (UVR) has potential protective, and confirmed detrimental effects on cancer risk. Several studies have examined recreational sun exposure in relation to breast cancer risk, and a recent meta-analysis found that moderate sun exposure (1–2 hrs/day) was associated with a decreased risk of breast cancer. Only one previous study in Canada has investigated UVR and breast cancer risk in an occupational setting.

Objectives The primary objective is to investigate the relationship between occupational solar UVR and breast cancer risk in Canada to inform on the dose-response relationship at the high end of the solar UVR spectrum.

Methods Questionnaire data from the Canadian Partnership for Tomorrow's Health (CanPath) will be utilized for participants in Ontario, Quebec, and Alberta. Information on sun exposure, longest job held, and breast cancer risk factors were obtained at baseline. Cancer outcomes have been ascertained via linkage with provincial cancer registries. A case-cohort approach was employed to facilitate job coding. Jobs codes were linked to a job exposure matrix (SUNJEM) to assign exposure. Breast cancer risk estimates (hazard ratios [HR]) and 95% confidence intervals (CI) will be estimated using a weighted Cox proportional hazard's regression with Prentice weights, controlling for potential confounders.

Results Preliminary results are available from the Ontario cohort, the analyses of additional cohorts are currently underway. From the underlying Ontario cohort 1,213 breast cancer cases met eligibility criteria. In the random sample of 2,500 women selected for the sub-cohort, the prevalence of occupational exposure (>2 hrs/day) to solar UVR was less than 2%. Preliminary HRs [0.995 (95%CI 0.684, 1.448)] suggest that occupational UVR does not have a protective or detrimental effect on the risk of breast cancer for Canadian women.

Conclusion This will be the second study to examine the relationship between occupational sun exposure and breast cancer risk in Canada, and the first to examine the dose-response relationship.

Cancer

O-19

COLORECTAL CANCER AMONG FARMERS IN THE AGRICAN COHORT STUDY¹Madar Talibov, Fabrice Morlais, Matthieu Meryet-Figuère, Mathilde Boulanger, Séverine Tual, Véronique Bouvier, Stephanie Perrier, Bénédicte Clin, Isabelle Baldi, Pierre Lebaillly. ¹INSERM U1086, France

10.1136/OEM-2021-EPI.16

Objective Specific farming types and tasks have never been studied in relation to colorectal cancer (CRC). We evaluated associations between 5 types of livestock and 13 types of crops in relation to CRC and its subsites within the Agriculture and Cancer (AGRICAN) study.

Methods AGRICAN cohort includes 181 842 agricultural workers living in 11 French geographical areas. Data on farming types and tasks was collected by self-administered questionnaires. We identified 2 609 CRC, 972 right colon, 689 left colon and 898 rectal incident cancer cases during follow-up from 2005 to 2015. Cox proportional hazards models were used to estimate hazard ratios (HR) and 95% confidence intervals (95% CI).

Results Significantly increased CRC risk was observed for farmers producing horses (HR=1.18, 95% CI 1.06–1.31), sunflower (HR=1.23, 95% CI 1.03–1.45) and field vegetables (HR=1.18, 95% CI 1.02–1.36). Positive associations were also observed for pig, poultry and wheat/barley. Some associations were observed only for specific subsites: left colon cancer was associated with fruit growing (HR=1.36, 95% CI 1.09–1.70) and potato (HR=1.28, 95% CI 1.05–1.57). Tasks related to livestock (animal care, insecticide treatment, disinfection of milking equipment and building) or to crop (haymaking, sowing, pesticide treatment, seed treatment, harvesting) were also associated with CRC. Duration and size of farming types/tasks increased the risk for some of the associations. Analysis stratified by gender suggested an interaction with several farming types/tasks.

Conclusion The current study showed original and positive findings for several farming types and tasks and CRC risk, overall and by subsites.

O-22

CANCER SURVEILLANCE AMONG PLASTICS AND RUBBER MANUFACTURING WORKERS IN ONTARIO, CANADA¹Nathan DeBono, Paul Demers, Chloë Logar-Henderson, Jill MacLeod, Mamadou Dakouo, Hunter Warden, Sharara Shakik. ¹International Agency for Research on Cancer, France

10.1136/OEM-2021-EPI.17

Objective Occupational exposure to agents used in plastics and rubber manufacturing have been associated with elevated risk of certain cancers. We sought to estimate cancer risk among workers with a history of employment in plastics and rubber manufacturing as part of an ongoing surveillance program in Ontario, Canada.

Methods The Occupational Disease Surveillance System (ODSS) cohort was established using workers' compensation claims data and includes 2.18 million workers employed between 1983–2014. Workers were followed for site-specific cancer diagnoses in the Ontario Cancer Registry through 2016. Cox-proportional hazard models were used to estimate adjusted hazard ratios (HR) and 95% confidence intervals (CI).

Results We identified 81,127 workers (69% male) ever-employed in plastics and rubber manufacturing industries or materials processing and product fabricating occupations. Compared to all other workers in the ODSS, workers in materials processing occupations had an elevated rate of lung cancer (HR 1.11, 95% CI: 1.02–1.20), which occurred almost exclusively among females (HR 1.38, 95% CI 1.20–1.58) in sex-stratified analyses. An elevated rate of breast cancer was observed among female labourers (HR 1.36, 95% CI: 1.01–1.82) and moulders (HR 1.47, 95% CI: 0.91–2.37) in plastics and rubber product fabricating occupations. Rates were elevated for esophageal, liver, stomach, prostate, and

kidney cancer in job-specific subgroups including mixing and blending, bonding and cementing, and labouring. Workers in the plastics product fabricating industry had modestly elevated rates of pancreatic and brain and nervous system cancer.

Conclusions Elevated rates of lung and breast cancer among females are consistent with other studies of women in plastics and rubber manufacturing and warrant further attention in Ontario. Results for digestive and other cancers are broadly consistent with exposure to known or suspected carcinogens in these industries and suggest new sites of potential concern.

0-272 EXPOSURE TO CARBAMATE INSECTICIDES AND RISKS OF NON-HODGKIN LYMPHOMAS IN THE FRENCH AGRICULTURE AND CANCER COHORT (AGRICAN)

¹Kenza Abdelmalki, Amandine Busson, Liacine Bouaoun, Séverine Tual, Alain Monnereau, Isabelle Baldi, Pierre Lebaillay, Kayo Togawa. ¹International Agency for Research on Cancer, France

10.1136/OEM-2021-EPI.18

Introduction Existing epidemiological studies have suggested a positive association between agricultural exposure to carbamates and risks of non-Hodgkin's lymphomas (NHL); however, the association remains inconclusive with most studies lacking statistical power to examine specific carbamates and subtypes of NHL.

Objectives We estimated the associations between carbamate insecticides and the risks of NHL overall and three major histological subtypes, multiple myeloma (MM), chronic lymphocytic leukaemia/small lymphocytic lymphoma (CLL-SLL), and diffuse large B-cell lymphoma (DLBCL) in the French prospective Agriculture and Cancer cohort (AGRICAN).

Methods At enrolment (2005–2007), participants completed a questionnaire on lifetime occupational history of agricultural practices, lifestyle habits, and medical history and were followed up until 2015 through linkage to the cancer registries. Information on pesticide use for different agricultural activities (crops, animals, barns, seeds) was crossed with the French crop-exposure matrix, PESTIMAT, enabling us to assess exposure to 19 specific carbamates (e.g., carbaryl, carbofuran) by type of activity. We estimated hazard ratios (HRs) with 95% confidence intervals (CIs) for NHL overall and its subtypes in relation to activity-specific exposure to carbamates (ever/never, duration) by fitting multivariate Cox proportional hazards models with age as the time scale. Non-users of pesticides were chosen as the reference group.

Results During the follow-up (median=8.9 years), 533 incident cases of NHL (MM=125, CLL-SLL=134, DLBCL=72) were diagnosed. Ever use of carbaryl on animals (HR=1.84, 95% CI=1.05–3.24) or barns (HR=1.86, 95% CI=1.06–3.27) were significantly associated with an increased risk of MM, but not for crops (HR=1.24, 95% CI=0.70–2.19). No significant trend by exposure duration were observed. None of the other carbamate insecticides were significantly associated with the risk of MM, CLL, DLBCL or NHL overall.

Conclusion Use of carbaryl on livestock production may lead to an increased risk of MM. Studies on activity-specific exposure pathways and levels are warranted to better understand the observed association.

0-276 DIESEL EXHAUST EXPOSURE AND SOMATIC MUTATIONS IN BLADDER TUMORS

¹Stella Koutros, Nicole Gonzalez, Nina Rao, Michael Dean, Donghyuk Lee, Amber Hurson, Dalsu Baris, Molly Schwenn, Ludmila Prokunina-Olsson, Melissa Friesen, Bin Zhu, Nathaniel Rothman, Debra Silverman, Alison Johnson. ¹U.S. National Cancer Institute, United States

10.1136/OEM-2021-EPI.19

Introduction Occupational diesel exhaust exposure has been linked to increased bladder cancer risk in epidemiologic studies, but few explore mechanistic links.

Objectives We examined the relationship between diesel exhaust exposure and somatic mutations and mutational signatures in bladder tumors.

Methods Targeted sequencing was conducted in bladder tumors from the New England Bladder Cancer Study (NEBCS), a population-based case-control study. Using data on 797 cases and 1,418 controls, two-stage polytomous logistic regression was used to calculate odds ratios (ORs) and 95% confidence intervals (CIs) to evaluate etiologic heterogeneity between bladder cancer subtypes and quantitative estimates of respirable elemental carbon. Poisson regression was used to evaluate associations between REC and mutational signatures including those from nitro-polycyclic aromatic hydrocarbons (nitro-PAHs) in the NEBCS and in publicly available whole-genome sequencing bladder cancer dataset from The Cancer Genome Atlas Project (TCGA) (N=412).

Results We observed significant heterogeneity in the diesel-bladder cancer risk relationship, with a strong positive association predominantly among cases with high-grade, non-muscle invasive TP53-mutated tumors compared to controls (ORTertile3vsUnexposed = 4.8, 95% CI=2.2,10.5; p-trend T mutations at CpG dinucleotides (Relative Risk (RR)=1.47, 95% CI=1.38,1.57). In muscle-invasive tumors, we also observed a positive association between diesel exposure and the diesel-associated nitro-PAH signatures of 1,6-dintropyrene (RR=1.93, 95% CI=1.28,2.92) and 3-nitrobenzoic acid (RR=1.97, 95% CI=1.33,2.92) in pooled analyses of NEBCS and TCGA.

Conclusion The relationship between diesel exhaust and bladder cancer was heterogeneous based on the presence of TP53 mutations in tumors. Assessing the impact of diesel exhaust exposure at CpG dinucleotides and any resultant transcriptional changes in bladder tumors may be valuable considering the wealth of data linking particulate exposures to altered DNA methylation. Future studies that can identify nitro-PAH signatures in tumors are of interest to support the findings generated here.

0-324 INCIDENCE OF POTENTIALLY EXPOSURE-RELATED CANCER SITES AMONG NORWEGIAN FIREFIGHTERS: 58 YEARS OF FOLLOW-UP

¹Niki Marjerrison, Jarle Jakobsen, Tom K Grimsrud, Johnni Hansen, Jan Ivar Martinsen, Karl-Christian Nordby, Raymond Olsen, Marit B Veierød, Kristina Kjaerheim. ¹Cancer Registry of Norway, Norway

10.1136/OEM-2021-EPI.20

Introduction Firefighters are exposed to a variety of known and suspected carcinogens through their work, and previous studies have found elevated risk for a number of cancer sites among firefighters.