Conclusions The impairment in respiration and hearing which is higher among the workers exposed to dust, fumes and noise at the workplace shows a need for future research to study if all safety measures are strictly adhered to. The impairment in near vision which was higher among the exposed workers needs detailed investigations into the cause and association with the working environment, if any.

P-100

MEDICAL ACCESSIBILITY AND UNDER-REPORTING OF OCCUPATIONAL DISEASES: EFFECT OF TRAVEL DISTANCE AND TRAVEL TIME

¹Ping Hui Chen, Pau Chung Chen. ¹National Taiwan University Hospital Hsinchu Branch, Taiwan

10.1136/OEM-2021-EPI.199

Introduction In Taiwan, outpatients' average travel distance (TD) is 17.68 km, and workers' TD and travel time (TT) for outpatient services are only 8.2 km and 27.6 mins. Poor medical accessibility of occupational outpatient service can also lead to under-reporting of occupational diseases (ODs).

Methods In Taiwan, Network of Occupational Diseases and Injuries Service (NODIS), composed of 9 major reporting hospitals, is an important surveillance system of ODs. Using NODIS's reporting data and manpower survey from 2008 to 2018, we calculate each town's incidence rate of occupation diseases (IROD) and expected IROD according to workers' occupations and job titles, and each town's shortest TD and TT to 9 major reporting hospitals is estimated by Google Maps' Distance Matrix API. Quasi-Poisson regression model is employed to investigate effect of TD and TT on IROD.

Results There are 8017 cases of suspected ODs in NODIS from 2008 to 2018, and 3306 cases are confirmed as definite ODs. Adjusted by workers' occupations and job titles, as TD/TT increases by 10 km/10 mins, IROD significantly decreases by 10.90%/10.74%, and less-disabled workers who have never stopped working or lost their jobs are more impeded by long TD and TT. Compared with towns with TD 45 km and offshore towns decreases by 38.93%, 39.58%, 50.03%, 55.01%, 65.71%, and 84.10%, and IRODs of towns with TT 10–15, 15–25, 25–35, 35–45, >45 mins and offshore towns decreases by 30.94%, 43.57%, 47.41%, 47.70%, 67.81%, and 85.29%. Around 40% ODs are under-reported due to poor medical accessibility.

Conclusion Our study shows how poor medical accessibility leads to serious under-reporting, and up to 40% ODs could be under-reported. Using this method, we can identify areas with poor medical accessibility and evaluate cost-effectiveness of adding reporting hospital.

P-101

OCCUPATIONAL DUST EXPOSURES AND CT FINDINGS OF INTERSTITIAL LUNG DISEASE AND CHRONIC OBSTRUCTIVE PULMONARY DISEASE

¹Inge Brosbøl Iversen, Kennet Sønderstgaard Thorup, Jesper Thygesen, Finn Rasmussen, Michael Brun Andersen, Elisabeth Bendstrup, Zara Ann Stokholm, Else Toft Würtz, Vivi Schlünssen, Jens Peder Ellekilde Bonde, Jakob Hjort Bønløkke, Hans Kromhout, Henrik Al. ¹Aarhus University Hospital, Denmark

10.1136/OEM-2021-EPI.200

Introduction Occupational dust exposure is associated with interstitial lung disease and chronic obstructive pulmonary disease, but little is known about the association with more discrete lung changes detected by lung scans.

Objectives To 1) analyze the relation between occupational dust levels and HRCT (high-resolution computed tomography) detected signs of pulmonary disease, and 2) map the prevalence of these signs in the Danish workforce.

Methods We are carrying out a cross-sectional study of 25,000 adults who underwent HRCT scans of the lungs 2011–2019 in Denmark. We will analyze the extent of emphysema and signs of pulmonary fibrosis such as, but not limited to, reticulation with Imbio Lung Texture Analysis™ of the HRCT scans. The DOC*X cohort provides annual information on occupation (ISCO-88) and industry for the total Danish workforce since 1976. Individual exposure levels are estimated using quantitative job exposure matrices for asbestos, crystalline silica, wood dust, and endotoxins. We will conduct adjusted analyses of exposure-response relations and tabulate distributions of emphysema and signs of pulmonary fibrosis for all occupations and industries.

Conclusion The study will provide new knowledge on pulmonary effects of current and past occupational dust levels. We will use a new software for objective identification and quantification of signs of pulmonary disease independent of diagnostic traditions. This sensitive and graduated measure of outcome will also enable more sensitive exposure-response analyses that include discrete signs of pulmonary disease. We expect this study to serve as a basis for targeted interventions of importance to the many that still have dusty work.

P-108

POPULATION-LEVEL ESTIMATES OF OCCUPATIONAL EXPOSURE TO CHLOROTHALONIL, 2,4-D, AND GLYPHOSATE IN CANADA'S AGRICULTURAL INDUSTRY (CAREX CANADA)

¹Ela Rydz, Cheryl Peters, Kristian Larsen. ¹Simon Fraser University, Canada

10.1136/OEM-2021-EPI.201

Introduction Certain pesticides may lead to adverse health outcomes including cancer; however, little is known about occupational pesticide exposure in Canada.

Objective The purpose of this study was to estimate the prevalence and likelihood of occupational exposure to chlorothalonil, 2,4-D, and glyphosate in Canada's agricultural industry.

Methods Lower and upper estimates were calculated using the Canadian Census of Population (CoP) and Census of Agriculture (CoA). We estimated the number of workers and proportion of farms applying 'herbicides' or 'fungicides' by farm type using CoA survey data. These values were multiplied to yield the number of workers at risk of exposure. Likelihood of exposure (exposed, probably exposed, possibly exposed) was qualitatively assigned using information on crop type, primary expected tasks, crop production practices, and residue transfer data. Agricultural workers who are at risk of exposure but were not captured by the CoA were identified using the CoP.

Results An estimated 37,700 to 55,800 workers (11–13% of agricultural workers) were exposed to glyphosate in Canada while 30,800 to 43,600 workers (9–11%) and 9,000 to 14,100 (3%) were exposed to 2,4-D and chlorothalonil,