Objectives The aim was to examine cancer incidence among Norwegian firefighters in sites with established associations to known carcinogenic occupational exposures. This included sites within the respiratory, urinary, and lympho-haematopoietic systems, as well as the skin and all sites combined.

Methods A newly established historical cohort of 3881 Norwegian firefighters who worked within the period 1950–2018 was linked to the Cancer Registry of Norway for incident cancers occurring in the period 1960–2018. We calculated standardized incidence ratios (SIR) with rates for the national male population as reference, and stratified SIR analyses by period of first employment, duration of employment, and time since first employment.

Results Elevated risk was seen for all sites combined (SIR 1.15, 95% confidence interval 1.07–1.23). Elevated risk of urinary tract cancer was observed among firefighters who began working before 1950, and with observation ≥ 40 years since first employment. With ≥ 40 years since first employment, risks of mesothelioma and laryngeal cancer were also elevated.

Conclusions Our patterns of increased incidence in cancer sites somewhat differs from observations in other studies, which may reflect that firefighting is a complex exposure that differs between countries, and that exposures have likely changed alongside changing fire contents, firefighting techniques, and equipment. However, our observed associations between firefighting and urinary tract cancer, laryngeal cancer, and mesothelioma have been observed in some studies previously, and may be related to carcinogenic occupational exposures. Differences in risk by period of employment potentially reflect improved quality and use of personal protective equipment, while stratification by time since first employment suggests that some cancers may have yet to develop among more recently employed firefighters.

0-392 EARLY DETECTION OF PROSTATE CANCER IN FIREFIGHTERS – A REGISTER-BASED STUDY OF PROGNOSTIC FACTORS AND SURVIVAL

¹Jarle Jakobsen, Marit B Veierød, Kristina Kjaerheim, Tom K Grimsrud, Sophie Dorothea Fosså, Bato Hammarström. ²Cancer Registry of Norway, Norway

10.1136/OEM-2021-EPI.21

Introduction Meta-analyses have shown firefighters to have increased prostate cancer (PCa) incidence compared to the general population. Firefighters are exposed to many occupational carcinogens, but no chemical agent is considered an established risk factor for PCa. Increased diagnostic intensity through regular health check-ups has been suggested as an explanation for increased PCa risk among firefighters.

Objectives To examine age at diagnosis, prognostic factors and survival of PCa in Norwegian firefighters and three other occupations undergoing occupational health check-ups, and comparing with PCa cases in the general population.

Methods All PCa cases diagnosed 1960–2017 were extracted from the Cancer Registry of Norway. Firefighters, military employees, pilots and police officers were identified through occupational data from Statistics Norway. Age at diagnosis, clinical stage, prostate-specific antigen (PSA), Gleason score, performance status and overall and PCa-specific survival in

cases in these occupations were compared with cases in the general population.

Results Firefighters were significantly younger at PCa diagnosis than cases in the general population in the periods 1960–1993 (mean difference 2.1 years) and 2007–2017 (mean difference 4.3 years). At diagnosis, firefighters had significantly lower PSA values, Gleason scores and performance status scores than the general population. Firefighters diagnosed 2007–2017 had lower risk of all-cause death compared to the general population (crude hazard ratio (HR) 0.71 (0.53–0.95)). No difference was found after adjusting for age at diagnosis (HR 1.03 (0.77–1.37)). Cases in the other three occupations generally also had lower age at diagnosis, better prognostic factors and better crude overall survival than cases in the general population.

Conclusion Younger age and better prognostic factors at PCa diagnosis among firefighters and other occupations with requirements for health check-ups compared with cases in the general population may indicate an increased diagnostic intensity, likely contributing to elevated PCa incidence in such occupations.

Cardiovascular

0-110 OCCUPATION AND INCIDENT STROKE IN A U.S. GENERAL POPULATION COHORT 45 YEARS OF AGE OR OLDER (REGARDS STUDY)

¹Tran Huynh, Igor Burstyn, Leslie MacDonald, Leslie McClure, Monika Safford, Virginia Howard. ²Drexel University, United States

10.1136/OEM-2021-EPI.22

Objective To identify occupational groups with greater risk of stroke within a cohort of older adults in the United States.

Methods Participants with occupational history data were selected from the REasons for Geographic and Racial Differences in Stroke (REGARDS) study (n=17,333) - a population-based prospective cohort of black and white men and women aged ≥ 45 years enrolled between 2003 and 2007. Self-reported occupational data were collected between 2011 and 2013. Participants were contacted every 6 months by telephone for self- or proxy-reported strokes later adjudicated by medical records through October 2018. Analyses focused on participants with a coded occupation who were employed in their longest held job prior to enrollment (n=15,016). Modified Poisson regression analyses were performed to estimate the relative risk (RR) for each two-digit Standard Occupational Classification (SOC) (2010 version) compared to all other occupation groups combined to highlight occupation groups at greatest risk of stroke.

Results We observed 735 strokes (5%) during 15 years of follow-up. Results showed that participants employed in 'food preparation and serving' (22 cases; 8%), 'sales' (69 cases; 7%), and 'production' occupations (80 cases; 6%) experienced a greater occurrence of stroke, compared with all other occupational groups combined. These associations were robust to adjustment for Framingham stroke risk score (that includes age, sex, smoking status), race, region, and BMI with adjusted RRs 1.5 (95% CI, 1.0, 2.3), 1.4 (95% CI, 1.1, 1.7) and 1.2 (95% CI, 1.0, 1.5), respectively. Findings may be biased by