Results Five hundred and twenty-seven participants were enrolled in the study with a mean age of 28.86 years and extremes ranging from 19 to 55 years. Prevalence of UL-MSDs in the study population was 10.6%. The univariate analysis showed that UL-MSDs in army officers were significantly associated with : age ($p < 10^{-3}$), job seniority ($p = 0.009$), weekly worked hours ($p = 0.007$), wearing a helmet ($p = 0.039$) and job strain ($p < 10^{-3}$). According to the multivariate analysis, determinants of the risk of UL-MSDs in the study population were: history of upper limbs trauma ($p = 0.002$, OR=3.1; CI 95% = [1.49; 6.44]), age ($p = 0.001$; OR=1.89; CI 95% = [1.30; 2.73]), occupational category ($p = 0.047$; OR= 0.78; CI 95% = [0.62; 0.99]) and irregular working hours ($p = 0.008$; OR=2.99; CI 95% = [1.32; 6.75]).

Conclusion Prevention of UL-MSDs in army officers represents major challenges for military health professionals. Keeping a good operational capacity of military personnel is dependent on the establishment of an effective global preventive approach that covers the various aspects of the work in this environment, while respecting its particularities.

P-244 FROM COMMERCIAL FISHERMEN TO RECREATIONAL BOATERS: FATALITIES AND LIFEJACKET USE INFORM THE NEED TO PROMOTE USE.

¹Laurel Kincl, Samantha Case, Viktor Bovbjerg, Amelia Vaughan. ¹Oregon State University – Corvallis, OR, United States

10.1136/OEM-2021-EPI.245

Objective Vessel disasters and falls overboard result in fatalities in the commercial fishing industry and recreational boating. We reviewed available fatality surveillance data to identify opportunities to promote lifejacket use in the US Pacific Northwest.

Methods Commercial fishing fatality information for 2000–2018 was obtained for all fatalities in Oregon and Washington waters from the Commercial Fishing Incident Database (CFID). Recreational boater fatality information was obtained from the Oregon and Washington State Marine Boards' publicly available information. Summary statistics were compiled and lifejacket policies reviewed for both occupational and recreational uses.

Results In Between 2000–2018 In Washington and Oregon there were 90 commercial fishing fatalities; only 5 (6%) victims were wearing a lifejacket, with 3 of those not properly worn. From 2000–2018 in Oregon there were 263 recreational boating fatalities with 76 (29%) victims wearing a lifejacket. In Washington, available data was limited from 2011–2017, there were 52 recreational boating fatalities with no lifejacket information. Commercial fishermen and recreational boaters over the age of 12 years are not required to wear lifejackets while boating, although Coast Guard-approved devices must be provided for each person onboard. Lifejacket use marketing promotions exist such as the 'Live to be Salty' campaign for commercial fishermen and 'Life Jackets and Seat Belts-It's Your Choice' for recreational boaters.

Conclusion The percent difference in lifejacket use between commercial fishermen and recreational boaters who suffered a fatality likely represents differences in the precipitating factors in the incidents, and the ease of wear for tasks. Primary prevention of vessel disasters and falls overboard is critical; as lifejackets are essential in the event of an emergency. A

regional intervention based on a successful program for lobstermen is proposed to bring lifejacket education, try-before-you-buy and discounts to commercial fishermen and recreational boaters that will be evaluated for impact and change of culture.

P-245 IMPACT OF FOOD ON DIVE'S SAFETY

¹Carla Torres, Antonio Saverghini, Simone Tavares. ¹McGill University, Canada

10.1136/OEM-2021-EPI.246

Introduction Professional diving is an activity that exposes the diver's body to numerous environmental and physiological challenges. These require adequate physical and mental health conditions as a prerequisite for the task's safe performance. The diver's diet is an underestimated risk factor that has not been properly studied so far. Thus, due to lack of dietary recommendations, many times divers make basic dietary mistakes that can lead to undesirable outcomes, from abdominal discomfort to death by drowning.

Objective To determine how food and physiological changes due to immersion and hyperbaric exposure can interfere with diving safety.

Methods

Literature review

Results Food-related factors can lead to life-threatening risk conditions, since situations that would normally be considered just inconvenient and uncomfortable can be potentially fatal when they occur underwater, such as vomiting, for example. Especially considering the diet specific factors: volume of the food, nutritional composition of the diet, and the time interval between the last meal and the dive, the food can be a hazard associated with risks that can compromise the safety in the dive. These risks can be divided into direct - such as hypoglycemia, dehydration, gastrointestinal barotraumas (mainly resulting from aerophagia and gastrointestinal producing conditions), and regurgitation with vomiting or bronchoaspiration - and indirect - such as increased cardiac risk and increased decompression stress (which could lead to decompression sickness despite the correct application of the decompression tables).

Conclusion Healthy eating is a fundamental part of ensuring a healthy lifestyle. Given the particularities imposed by immersion and hyperbaric exposure, the diver's diet should be considered an important aspect in promoting health and diving safety. Awareness of the risks and knowledge of measures to increase the safety of diving can lead to beneficial changes in habits and, consequently, in the safety of diving operations.

P-246 'HEALTH AND WORKING CONDITIONS OF WASTE PICKERS: SCOPING REVIEW.'

¹Lina Paola Escobar Rincón, Oneys De Arco Canoles. ¹Universidad Nacional de Colombia, Colombia

10.1136/OEM-2021-EPI.247

Introduction It is estimated that between 15 to 20 million people in the world recover materials from waste in an informal condition. Many living in poverty, become involved in the activity because they do not have an accessible job opportunity. This need leads them to be exposed to unhealthy,