

Original research

# Military occupation and testicular germ cell tumour risk among US Air Force servicemen

Hristina Denic-Roberts <sup>1,2</sup>, Katherine McGlynn,<sup>3</sup> Jongeun Rhee,<sup>3</sup> Celia Byrne,<sup>1</sup> Michael Lang,<sup>4</sup> Paul Vu,<sup>5</sup> Mark Purdue,<sup>3</sup> Jennifer Rusiecki<sup>1</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2022-108628>).

<sup>1</sup>Department of Preventive Medicine and Biostatistics, Uniformed Services University, Bethesda, Maryland, USA

<sup>2</sup>Henry M Jackson Foundation for the Advancement of Military Medicine, Bethesda, Maryland, USA

<sup>3</sup>Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland, USA

<sup>4</sup>Epidemiology Consult Service, US Air Force School of Aerospace Medicine, Wright-Patterson AFB, Ohio, USA

<sup>5</sup>US Air Force Medical Readiness Agency, US Air Force Medical Service, Falls Church, Virginia, USA

## Correspondence to

Dr Jennifer Rusiecki, Department of Preventive Medicine and Biostatistics, Uniformed Services University, Bethesda, MD 20814, USA; [jennifer.rusiecki@usuhs.edu](mailto:jennifer.rusiecki@usuhs.edu)

Received 26 August 2022  
Accepted 21 February 2023  
Published Online First  
27 March 2023



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Denic-Roberts H, McGlynn K, Rhee J, et al. *Occup Environ Med* 2023;**80**:312–318.

## ABSTRACT

**Objectives** Testicular germ cell tumours (TGCTs) are the most commonly diagnosed malignancy among active duty US military servicemen. Occupational risk factors may play a role in TGCT aetiology, although the evidence is inconclusive. The objective of our study was to investigate associations between military occupations and TGCT risk among US Air Force (USAF) servicemen.

**Methods** This nested case–control study among active duty USAF servicemen obtained information on military occupations for 530 histologically confirmed TGCT cases diagnosed during 1990–2018 and 530 individually matched controls. We determined military occupations using Air Force Specialty Codes ascertained at two time points: at case diagnosis and at a time point on average 6 years earlier. We computed adjusted ORs and 95% CIs from conditional logistic regression models to evaluate associations between occupations and TGCT risk.

**Results** The mean age at TGCT diagnosis was 30 years. Increased TGCT risk was observed for pilots (OR=2.84, 95% CI: 1.20–6.74) and servicemen with aircraft maintenance jobs (OR=1.85, 95% CI: 1.03–3.31) who held those jobs at both time points. Fighter pilots (n=18) and servicemen with firefighting jobs (n=18) at the time of case diagnosis had suggestively elevated TGCT odds (OR=2.73, 95% CI: 0.96–7.72 and OR=1.94, 95% CI: 0.72–5.20, respectively).

**Conclusions** In this matched, nested case–control study of young active duty USAF servicemen, we found that pilots and men with aircraft maintenance jobs had elevated TGCT risk. Further research is needed to elucidate specific occupational exposures underlying these associations.

## INTRODUCTION

The incidence of testicular germ cell tumours (TGCTs), the most common (98%) type of testicular cancer, has increased in the USA and other developed countries over the past several decades.<sup>1</sup> For US men aged 15–44 years,<sup>2,3</sup> as well as for all active duty US military servicemen,<sup>4</sup> TGCT is the most commonly diagnosed malignancy. While known to be associated with other male reproductive disorders, such as cryptorchidism, hypospadias and impaired fertility,<sup>5</sup> the aetiology of TGCT is not well understood. Established TGCT risk factors include European ancestry, personal or family TGCT history, and taller adult height.<sup>6</sup> Events in utero almost certainly influence TGCT risk<sup>7</sup>; however, recent evidence suggests that adolescent and adult risk factors may also play an important

## WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Occupations including firefighting, aircraft maintenance and being a pilot have been associated with increased risk of testicular germ cell tumours (TGCTs), the most common type of testicular cancer, although the evidence is inconclusive, especially among the US military service members.
- ⇒ No studies to date have comprehensively investigated associations between various types of jobs performed during military service and TGCT risk.

## WHAT THIS STUDY ADDS

- ⇒ Our matched case–control study conducted among young active duty US Air Force servicemen observed that pilot and aircraft maintenance occupations were associated with elevated TGCT risk.
- ⇒ Fighter pilots and firefighters also had suggestively elevated TGCT risk, while the majority of other occupations and jobs we examined were not associated with increased risk.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Further research, including a comprehensive study of military personnel from all service branches, is needed to elucidate specific occupational exposures underlying the associations we observed.

role in TGCT aetiology.<sup>8</sup> Certain occupations have been associated with increased TGCT risk, including firefighting,<sup>9–10</sup> aircraft maintenance among both military<sup>11–12</sup> and civilian populations,<sup>13</sup> as well as service as a military pilot and aircrew,<sup>14–15</sup> and as a fighter pilot.<sup>16</sup> Despite inconsistencies, these occupational associations led to hypotheses that job-related chemical exposures, such as to perfluoroalkyl and polyfluoroalkyl substances (PFAS),<sup>17–19</sup> solvents, paints, and hydrocarbons in degreasing/lubricating agents and lubricating oils (eg, methylcholantene),<sup>11–12</sup> may increase TGCT risk, although confounding from factors related to socioeconomic status (SES) may also play a role.<sup>15–16</sup>

The predominantly young and male US military population, where military personnel may experience different occupational exposures throughout

their career/deployments, is valuable for studying occupational risk factors for TGCT. Prior epidemiological studies reported elevated risk of TGCT in military pilots and aircrew,<sup>15</sup> and fighter pilots<sup>16</sup>; however, to our knowledge, no peer-reviewed studies to date have investigated associations of different types of jobs performed during military service with TGCT risk. To address this research gap, the primary aim of the present study was to investigate associations between military occupations and TGCT risk among active duty US Air Force (USAF) servicemen. In a secondary aim, we additionally explored these associations by tumour morphology to measure histology-specific risks.

## METHODS

### Study population and outcome ascertainment

The present study is derived from a matched case–control study investigating the association between serum PFAS concentrations and TGCT risk, in which military occupation at different time points was ascertained.<sup>20</sup> Our study population was comprised of active duty USAF male personnel who had serum samples stored at the Department of Defense Serum Repository (DoDSR), a central repository maintained by the Armed Forces Health Surveillance Division (AFHSD) of the Defense Health Agency's Public Health Directorate.<sup>21</sup> The DoDSR blood samples are drawn from all military personnel at the time of joining the military (“military accession”) and, on average, every 2 years thereafter for mandatory HIV testing.<sup>22</sup> Currently, the DoDSR stores more than 60 million serial serum samples at  $-30^{\circ}\text{C}$  from more than 10 million active duty and reserve personnel in Silver Spring, Maryland. Eligible cases were identified through data linkage between the DoDSR and the Department of Defense (DoD) Automated Cancer Tumor Registry that records patients diagnosed with and/or treated for cancer at military treatment facilities. Through data linkage, we identified 530 incident TGCT cases during 1990–2018 (Third Edition of the International Classification of Diseases for Oncology site code C62 with classic seminoma morphology codes 9060–9062 and 9064 and non-seminoma morphology codes 9065–9102) among active duty USAF personnel with at least one banked DoDSR pre-diagnostic serum sample. TGCT cases with a history of any other malignancies before TGCT diagnosis date, except for non-melanoma skin cancer, were excluded from our study. Using risk set sampling with replacement, 530 controls were individually matched to each case on race/ethnicity, year of military accession ( $\pm 1$  year), date of birth ( $\pm 1$  year) and the collection date(s) of the case's serum sample(s) selected for study ( $\pm 1$  year). Each control was randomly selected from a population of active duty USAF servicemen with samples in the DoDSR who had no history of any malignancies before their matched case's TGCT diagnosis date, except for non-melanoma skin cancer. No control became a case during the study period.

Demographic and military characteristics, including occupational data, for all cases and controls were obtained from the Defense Medical Surveillance System (DMSS), the AFHSD, Defense Health Agency, Silver Spring, Maryland (1986–2018). This study did not include any interviews.

### Exposure ascertainment

Air Force occupation was ascertained from Air Force Specialty Codes (AFSCs) obtained from the DMSS for each serviceman in our study. These alphanumeric job codes classify USAF military personnel into specific occupations/career fields, hereafter referred to as *occupations*. Each AFSC corresponds to a job title specific to enlisted personnel or officers. Guided by the

Air Force Enlisted Classification Directory (AFECD)<sup>23</sup> and the Air Force Officer Classification Directory (AFOCD),<sup>24</sup> we first classified each airman's occupation based on career group and field specifically indicated by the characters within the AFSC. Because people in the same career field/group may have a range of different jobs and responsibilities, we further grouped participants into occupational groups based on job descriptions (in the AFECD/AFOCD) and professional expertise (PV and ML) about similar job exposures. This process resulted in some occupational groups including both enlisted personnel and officers (eg, administrative/support jobs). Online supplemental table 1 shows a listing of all job titles in each occupation/occupational group.

We ascertained occupational information at two time points: at the time of serum draw (time 1) and at the case's TGCT diagnosis date (time 2). Because occupational information at time 1 was ascertained shortly after military accession, on average 2.4 years post-accession, a large proportion (45%) of our study population was not occupationally qualified (eg, basic enlisted airman) at that time. We, therefore, primarily relied on occupational information at time 2, on average 8.2 years post-accession, when approximately only 3% of the population was non-occupationally qualified. For selected occupations of interest based on prior literature (ie, pilot, fighter pilot, firefighting and aircraft maintenance),<sup>8 16 25–27</sup> we additionally created occupational groupings that indicated being in a given occupation at both time points (eg, pilot at time 1 and at time 2, aircraft maintenance at time 1 and at time 2). We did not have information on occupation timing and duration (eg, when a serviceman started or ended a job).

### Statistical analyses

We first compared distributions of the demographic and military characteristics between cases and controls. To examine associations between different occupations/occupational groups and TGCT risk, we used conditional logistic regression models conditioned on the individually matched case–control pairs. We estimated adjusted ORs (aORs) and 95% CIs comparing risk of TGCT among active duty USAF servicemen in the select occupation with the risk of TGCT among those servicemen in all other occupations. All of the conditional logistic regression models were adjusted for number of deployments before the case diagnosis year (0, 1, 2+). Occupational groups consisting of both enlisted personnel and officers (eg, administrative/support) were further adjusted for the service member being either an officer or enlisted (ie, grade) at case diagnosis. With an aim of accounting for grade, for pilot-related occupations of interest (comprised entirely of officers), we also compared TGCT risk between officer pilots and a reference group of officer non-pilots.

To test whether associations between occupation and TGCT risk were specific to either seminomas or non-seminomas, we performed conditional logistic regression analyses stratified by morphology of the cases. We conducted case-only analyses of TGCT subtypes using unconditional logistic regression adjusting for age at diagnosis, race/ethnicity, number of deployments before TGCT diagnosis and grade, where appropriate, in order to statistically test for differences in associations by morphology.<sup>28</sup> We calculated p values to test for OR homogeneity across case morphology.

### Sensitivity analyses

With an aim of excluding occupational exposures that occurred too close to the TGCT diagnosis to have aetiological relevance, we performed a sensitivity analysis excluding cases and controls with fewer than 3 years of service ( $n=202$ ) at the time of the case diagnosis date. We additionally performed a sensitivity analysis

among a subset of participants with information on height ( $n=689$ ) in order to assess the potential confounding effect of height on the main associations. We compared conditional logistic regression estimates adjusted for number of deployments before case diagnosis and grade, where appropriate, with the estimates additionally adjusted for height (in centimetres) at military accession. To assess the sensitivity of our findings, we also performed the main analyses without including the number of deployments as a potential confounding variable, but the results were largely unchanged (data not shown).

For each analysis, we selected occupations/occupational groups for which there were at least five cases and five controls. All of the analyses were performed in SAS V.9.4 (SAS Institute).

## RESULTS

**Table 1** shows characteristics of cases and controls. Our study population joined the military between 1986 and 2017 (median year 1998). TGCT cases were diagnosed between 1990 and 2018 (median year 2007) at an average age of 29.8 years. Slightly more than half of the cases (55.3%) were diagnosed with seminomas. Cases and matched controls were predominantly of non-Hispanic white background (79.6%), while 10.8% identified as Hispanic, and 2.4% as non-Hispanic black. At the time of case diagnosis, a higher proportion of cases (22.6%) than controls (17.9%) were officers. Although a large proportion of height and weight information at military accession was missing for both cases and controls (~35%), the median height and weight were similar between the two groups. A higher proportion of cases (72.8%) than controls (67.4%) had not deployed before the case diagnosis year. On average, cases and controls had served in the USAF for 8.2 years at the time of case diagnosis. The average time between the serum sample collection and case diagnosis was 5.8 years.

Findings from analyses of occupation at the time of case diagnosis and odds of TGCT are presented in **table 2**. Pilots (including pilot trainees) had elevated TGCT odds (aOR=1.78, 95% CI: 1.06–3.01). The association was stronger, though not statistically significant, for fighter pilots (aOR=2.73, 95% CI: 0.96–7.72). Airmen with aircraft maintenance jobs also had elevated TGCT odds (aOR=1.54, 95% CI: 1.04–2.29). Fire protection occupation (ie, firefighting) was associated with non-significantly increased TGCT odds (aOR=1.94, 95% CI: 0.72–5.20), while servicemen with occupations in communication/electronics/wire system maintenance, aerospace medical/surgical service, intelligence and aircraft crew had non-significantly reduced TGCT odds (aOR range: 0.52–0.67).

In analyses stratified by tumour morphology (**table 3**), we generally observed similar occupation findings for each subtype, although associations with the command and control systems operations and aerospace maintenance occupations were significantly stronger for non-seminomas than seminomas (case-only analysis  $p=0.04$  for each occupation). The association between aircraft maintenance jobs and non-seminomas was also slightly stronger (aOR=2.00, 95% CI: 1.13–3.52) than for seminomas (aOR=1.28, 95% CI: 0.72–2.27, case-only analysis  $p=0.06$ ).

**Table 4** shows associations with TGCT risk for selected occupational groups of interest at the time of serum collection and case diagnosis. The aOR for being a pilot at the time of serum collection was 2.54 (95% CI: 1.11–5.79) and at case diagnosis, it was 1.72 (95% CI: 0.98–3.02). Those who were pilots at both time points had the highest TGCT risk (aOR=2.84, 95% CI: 1.20–6.74). We observed a similar pattern of associations, although lower in magnitude, for a combined occupation

**Table 1** Select characteristics of the US Air Force study population

Characteristic	TGCT cases (N=530)		Controls (N=530)	
	N (%)		N (%)	
Age at serum sample collection (year)				
<20	141 (26.6)		134 (25.3)	
20–24	194 (36.6)		220 (41.5)	
25–29	95 (17.9)		76 (14.3)	
30–34	64 (12.1)		66 (12.4)	
35–39	28 (5.3)		27 (5.1)	
40+	8 (1.5)		7 (1.3)	
Age at TGCT diagnosis (year)				
<20	10 (1.9)			
20–24	111 (20.9)			
25–29	159 (30)			
30–34	113 (21.3)			
35–39	89 (16.8)		–	
40+	48 (9.1)		–	
Mean±SD	29.8±6.6		–	
Race/ethnicity at case diagnosis				
Non-Hispanic white	422 (79.6)		422 (79.6)	
Non-Hispanic black	13 (2.4)		13 (2.4)	
Hispanic	57 (10.8)		57 (10.8)	
Asian/Pacific Islander	6 (1.1)		6 (1.1)	
Native American/Alaska Native	5 (0.9)		5 (0.9)	
Other	9 (1.7)		9 (1.7)	
Unknown	18 (3.4)		18 (3.4)	
Education at case diagnosis				
≤High school	261 (49.3)		268 (50.6)	
Some college	120 (22.6)		138 (26)	
Bachelor's	80 (15.1)		57 (10.8)	
Advanced degree	68 (12.8)		66 (12.4)	
Unknown	1 (0.2)		1 (0.2)	
Grade at case diagnosis				
Enlisted (E01–E10)	410 (77.4)		435 (82.1)	
Officer (O1–O10)	120 (22.6)		95 (17.9)	
Height (m) at military accession				
Mean±SD	1.65±0.24		1.61±0.21	
% Missing	190 (35.8)		181 (34.2)	
Weight (kg) at military accession				
Mean±SD	69.9±2.59		69.5±2.48	
% Missing	190 (35.8)		181 (34.2)	
Number of deployments before case dx year				
0	386 (72.8)		354 (67.4)	
1	81 (15.3)		92 (17.4)	
2	35 (6.6)		50 (9.4)	
≥3	28 (5.3)		32 (5.8)	
Years of service at serum sample collection				
Mean±SD	2.4±3.3		2.4±3.3	
Years of service at case dx				
Mean±SD	8.2±5.3		8.2±5.2	
Time between serum collection and case dx (years)				
Mean±SD	5.8±4.4		5.8±4.4	
Calendar year of military accession				
Median (range)	1998 (1986–2017)		1998 (1986–2017)	
Calendar year of serum sample draw				
Median (range)	1999 (1988–2017)		1999 (1989–2017)	

continued



Table 1 continued

Characteristic	TGCT cases (N=530)	Controls (N=530)
Calendar year at case dx		
Median (range)	2007 (1990–2018)	2007 (1990–2018)
Tumour morphology		
Seminoma	293 (55.3)	
Non-seminoma	237 (44.7)	
Controls individually matched to cases on: race/ethnicity, military accession year ( $\pm 1$ year), date of birth ( $\pm 1$ year) and serum sample collection date ( $\pm 1$ year). *Defined as a deployment record with a length of $\geq 31$ days outside of the USA. dx, diagnosis; TGCT, testicular germ cell tumour.		

of pilots and pilot trainees. There were too few fighter pilots and firefighters in the study to conduct the analysis at both time points. Having an aircraft maintenance job at case diagnosis was significantly associated with TGCT (aOR=1.54, 95% CI: 1.04–2.29); the elevated risk was stronger for those in aircraft maintenance at both time points (aOR=1.85, 95% CI: 1.03–3.31).

Because being a pilot or a pilot trainee is a job specific to officers, we could not adjust the main case–control models for grade. However, in addition to comparing the TGCT risk between pilots and all non-pilots (ie, a reference group consisting of both enlisted personnel and non-pilot officers), we compared pilot officers separately to all enlisted personnel and to all other officers who were not pilots (online supplemental table 2). For these comparisons, we used pilot and pilot trainee occupation at both time points. Compared with all enlisted personnel, pilot officers (excluding trainees) had a significantly elevated TGCT risk (aOR=2.90, 95% CI: 1.22–6.89). This association slightly attenuated when comparing pilots with the rest of the non-pilot officers (aOR=2.43, 95% CI: 0.96–6.13). We observed a similar pattern when combining pilots and pilot trainees (online supplemental table 2).

Results from the sensitivity analysis excluding 202 servicemen with less than 3 years of service at the time of case diagnosis (online supplemental table 3) were generally similar to those in the main analyses among the full case–control population presented in table 2. The pilot occupation (excluding or including pilot trainees) remained significantly associated with elevated TGCT risk (aOR<sub>pilot</sub>=1.80, 95% CI: 1.02–3.18 and aOR<sub>pilot/pilot trainee</sub>=1.76, 95% CI: 1.01–3.05), while the association with aircraft maintenance-related jobs slightly attenuated (aOR=1.46, 95% CI: 0.94–2.27). Compared with the main analyses of the full case–control population, the associations between some occupations/occupational groups and TGCT risk strengthened in the sensitivity analysis but were not statistically significant (eg, aOR=1.83, 95% CI: 0.80–4.15 for command and control systems operations; aOR=2.04, 95% CI: 0.84–4.96 for jobs involving fuels/petroleum). Sensitivity analyses excluding 371 servicemen without information on height are shown in online supplemental table 4. Further adjustment for height did not significantly alter the associations (ie, the point estimates changed by <10%).

## DISCUSSION

In this nested case–control study of USAF servicemen, we found significantly increased risks of TGCT among pilots and aircraft maintenance workers, while non-significantly elevated risks were observed for the subcategory of fighter pilots and for firefighters. We also observed suggestions of reduced TGCT risk among airmen with occupations/occupational groups in

Table 2 Case–control associations between occupation at case diagnosis and TGCT

	Cases (N=530)	Controls (N=530)	OR (95% CI)
<b>Occupation (AFSC)</b>	<b>N</b>	<b>N</b>	<b>OR (95% CI)</b>
Aircrew operations (1A)	14	23	0.67 (0.34–1.35)
Command and control systems operations (1C)	17	14	1.22 (0.60–2.49)
Aerospace maintenance (2A)	103	93	1.12 (0.82–1.54)
Fuels (2F)	11	9	1.32 (0.54–3.22)
Material management (2S)	11	11	1.01 (0.43–2.35)
Transportation (2T)	20	23	0.93 (0.50–1.73)
Munitions and weapons (2W)	27	34	0.77 (0.45–1.32)
Communication-electronics/wire systems maintenance (2E)	12	22	0.56 (0.27–1.13)
Communications/network (3C)	11	10	1.07 (0.43–2.64)
Cyberspace support (3D)	23	18	1.35 (0.70–2.61)
Security forces (3P)	39	38	1.08 (0.67–1.73)
Fire protection (3E7×1)	12	6	1.94 (0.72–5.20)
Aerospace medical/surgical service (4N)	9	17	0.52 (0.23–1.17)
Pilot (11)	36	23	1.72 (0.98–3.02)
Fixed-wing pilot (11A, 11K, 11M, 11R, 11S, 11T)	20	17	1.22 (0.63–2.39)
Fighter pilot (11F)	13	5	2.73 (0.96–7.72)
Pilot trainee (92T0)	10	5	1.77 (0.53–5.88)
Pilot/pilot trainee (11, 92T0)	46	28	<b>1.78 (1.06–3.01)</b>
Space, nuclear and missile (13)	10	12	0.89 (0.38–2.07)
Civil engineer (3E, 32E)*	38	36	1.07 (0.67–1.72)
Intelligence (1N, 14N)*	19	30	0.61 (0.34–1.11)
<b>Occupational group based on job description (AFSC)</b>			
Aircraft maintenance (2A, 21A)*	69	47	<b>1.54 (1.04–2.29)</b>
Aircraft crew (12, 13, 1A, 2A)*	24	40	0.60 (0.35–1.04)
Ammunition (3E, 2W)	30	34	0.88 (0.53–1.46)
Fuels/petroleum (2A, 2F, 63)*	15	12	1.42 (0.66–3.08)
Electrical/electronics (2A, 2E, 2P, 3D, 3E)	22	32	0.62 (0.34–1.12)
Mechanic/other maintenance (2A, 2E, 2M, 2T, 3E, 4A)	21	29	0.65 (0.36–1.20)
Law enforcement/security (3P, 31P, 81)*	42	39	1.20 (0.76–1.91)
Radio/communications/radar (1C, 2A, 2E, 3C, 3D, 33S)*	24	23	1.04 (0.57–1.87)
Medical*†	35	31	1.13 (0.69–1.94)
Administrative/support*‡	139	137	0.99 (0.75–1.33)
Miscellaneous/other/unknown*§	44	56	0.78 (0.50–1.20)
All conditional logistic regression models adjusted for number of deployments before case diagnosis (0, 1, 2+).			
Bold indicative of statistical significance.			
Comparison (non-exposed) group is everyone not in selected occupation/occupational group.			
*Models additionally adjusted for grade at case diagnosis.			
†Medical AFSCs: 42, 44, 45, 46, 47, 48, 4C, 4D, 4F, 4N, 4P, 4R, 4Y, 92M1.			
‡Administrative/support AFSCs: 13, 16, 17, 18, 20, 21, 27, 43, 51, 60, 62, 63, 64, 65, 97, 10C, 14N, 1C, 1N, 1W, 21R, 2E, 2G, 2R, 2S, 2T, 32E, 33S, 34M, 36M, 36P, 38F, 38P, 3A, 3C, 3D, 3M, 3N, 3S, 3V, 41A, 4A, 4E, 52R, 5R, 6C, 6F, 7S, 8R, 8M000, 9E.			
§Miscellaneous/other/unknown AFSCs: 45, 61, 62, 15W, 1C, 1P, 1S, 1T, 1U, 2A, 2E, 2T, 2W, 3E, 4B, 8F, 9S, 90G0, 9T000.			
AFSC, Air Force Specialty Code; TGCT, testicular germ cell tumour.			

communication/electronics/wire system maintenance, aerospace medical/surgical service, intelligence and aircraft crew.

One of our most consistent findings of elevated TGCT risk for pilots is comparable with the findings of some of the previous studies conducted among USAF servicemen,<sup>14–16</sup> although not all.<sup>29</sup> Previous

**Table 3** Case-control associations between occupation at case diagnosis and TGCT, stratified by tumour morphology

Occupation (AFSC)	Seminoma (293 case-control pairs)			Non-seminoma (237 case-control pairs)			P value*
	N case	N control	OR (95% CI)	N case	N control	OR (95% CI)	
Aircrew operations (1A)	8	12	0.66 (0.26–1.72)	6	11	0.75 (0.27–2.13)	0.86
Command and control systems operations (1C)	6	8	0.74 (0.26–2.14)	11	6	1.88 (0.68–5.19)	<b>0.04</b>
Aerospace maintenance (2A)	43	49	0.87 (0.56–1.35)	60	44	1.56 (0.97–2.50)	<b>0.04</b>
Fuels (2F)	8	3	–	3	6	–	–
Material management (2S)	7	5	1.33 (0.42–4.22)	4	6	–	–
Transportation (2T)	10	12	0.85 (0.35–2.07)	10	11	1.01 (0.42–2.42)	0.74
Munitions and weapons (2W)	13	18	0.74 (0.35–1.55)	14	16	0.79 (0.36–1.74)	0.95
Communication-electronics/wire systems maintenance (2E)	6	12	0.50 (0.19–1.33)	6	10	0.69 (0.25–1.93)	0.90
Communications/network (3C)	4	2	–	7	8	0.83 (0.28–2.47)	–
Cyberspace support (3D)	12	9	1.40 (0.59–3.35)	11	9	1.21 (0.44–3.34)	0.89
Security forces (3P)	24	18	1.39 (0.72–2.65)	15	20	0.75 (0.37–1.55)	0.08
Fire protection (3E7×1)	3	4	–	9	2	–	–
Aerospace medical/surgical service (4N)	6	8	0.72 (0.25–2.10)	3	9	–	–
Pilot (11)	24	15	1.69 (0.87–3.31)	12	8	2.09 (0.73–6.01)	0.75
Fixed-wing pilot (11A, 11K, 11M, 11R, 11S, 11T)	12	10	1.21 (0.52–2.80)	8	7	1.40 (0.45–4.35)	0.70
Fighter pilot (11F)	9	4	–	4	1	–	–
Pilot trainee (92T0)	5	4	–	5	1	–	–
Pilot/pilot trainee (11, 92T0)	29	19	1.62 (0.87–3.03)	17	9	2.60 (0.96–7.03)	0.62
Space, nuclear and missile (13)	6	8	0.78 (0.27–2.27)	4	4	–	–
Civil engineer (3E, 32E)†	19	23	0.80 (0.42–1.49)	19	13	1.59 (0.75–3.39)	0.89
Intelligence (1N, 14N)†	13	14	0.90 (0.42–1.92)	6	16	0.35 (0.12–0.97)	0.24
<b>Occupational group based on job description (AFSC)</b>							
Aircraft maintenance (2A, 21A)†	29	23	1.28 (0.72–2.27)	40	24	<b>2.00 (1.13–3.52)</b>	0.06
Aircraft crew (12, 13, 1A, 2A)†	13	22	0.57 (0.27–1.19)	11	18	0.57 (0.24–1.34)	0.60
Ammunition (3E, 2W)	13	17	0.78 (0.37–1.67)	17	17	0.96 (0.47–1.95)	0.37
Fuels/petroleum (2A, 2F, 63)†	12	4	–	3	8	–	–
Electrical/electronics (2A, 2E, 2P, 3D, 3E)	12	20	0.53 (0.23–1.18)	10	12	0.78 (0.32–1.91)	0.67
Mechanic/other maintenance (2A, 2E, 2M, 2T, 3E, 4A)	9	16	0.49 (0.20–1.21)	12	13	0.79 (0.34–1.86)	0.23
Law enforcement/security (3P, 31P, 81)†	25	19	1.41 (0.74–2.66)	17	20	0.94 (0.47–1.89)	0.12
Radio/communications/radar (1C, 2A, 2E, 3C, 3D, 33S)†	11	12	0.94 (0.41–2.15)	13	11	1.10 (0.47–2.59)	0.30
Medical‡	25	19	1.32 (0.66–2.64)	12	13	0.90 (0.37–2.18)	0.61
Administrative/support‡§	89	76	1.26 (0.85–1.86)	50	61	0.77 (0.50–1.20)	0.18
Miscellaneous/other/unknown†¶	20	33	0.58 (0.32–1.07)	24	23	1.02 (0.53–1.95)	0.46

All conditional logistic regression models adjusted for number of deployments before case diagnosis (0, 1, 2+).

Bold indicative of statistical significance.

Comparison (non-exposed) group is everyone not in selected occupation/occupational group.

\*P value from tests of OR homogeneity across case morphology from case-only unconditional logistic regression models adjusting for age at diagnosis, race/ethnicity, number of deployments before diagnosis and grade where appropriate.

†Models additionally adjusted for grade at case diagnosis.

‡Medical AFSCs: 42, 44, 45, 46, 47, 48, 4C, 4D, 4F, 4N, 4P, 4R, 4Y, 92M1.

§Administrative/support AFSCs: 13, 16, 17, 18, 20, 21, 27, 43, 51, 60, 62, 63, 64, 65, 97, 10C, 14N, 1C, 1N, 1W, 21R, 2E, 2G, 2R, 2S, 2T, 32E, 33S, 34M, 36M, 36P, 38F, 38P, 3A, 3C, 3D, 3M, 3N, 3S, 3V, 41A, 4A, 4E, 52R, 5R, 6C, 6F, 7S, 8R, 8M000, 9E.

¶Miscellaneous/other/unknown AFSCs: 45, 61, 62, 15W, 1C, 1P, 1S, 1T, 1U, 2A, 2E, 2T, 2W, 3E, 4B, 8F, 9S, 90G0, 9T000.

AFSC, Air Force Specialty Code; TGCT, testicular germ cell tumour.

studies that found elevated risk/odds of TGCT among aviators slightly differed with respect to the exposure metrics. In one study conducted among active duty USAF officers who served for at least a year during 1975–1989, the age-adjusted incidence rate of testicular cancer was significantly elevated among those with a professional history of flying compared with those with no flying history (rate ratio (RR)=1.84, 99% CI: 1.19–2.86).<sup>14</sup> Similarly, in a study of white active duty USAF officers who were admitted to US military treatment facilities during 1988–1999, those who flew for at least 1 hour had an elevated risk of testicular cancer (OR=1.74, 95% CI: 1.04–2.92), and there was a suggestion of a dose–response relationship with flight hours.<sup>15</sup> In a more recent study conducted among active duty USAF officers who entered the military during 1970–2004,

fighter aviators (including pilots and backseat aircrew) with at least 100 hours of flying on a fighter airframe had increased age-adjusted and race-adjusted incidence of testicular cancer (OR=1.29, 95% CI: 1.15–2.12).<sup>16</sup> However, in a study of active duty USAF officers who entered the military during 1986–2006, Robbins *et al* did not find an elevated incidence rate of testicular cancer comparing fighter pilots with all other officers (incidence rate ratio (IRR)=0.92, 95% CI: 0.56–1.52).<sup>29</sup> Those investigators relied on ‘usual occupation’ for fighter pilots based on the length of time spent in the occupation and, therefore, approximately 8% of the comparison group also included officers who were fighter pilots at some point in their career.<sup>29</sup> This exposure misclassification, and a shorter study follow-up period, could have contributed to the null finding for testicular cancer risk.

**Table 4** Case–control associations between select occupations at two time points and TGCT risk

Occupation/occupational group (AFSC)	Cases (N=530)	Controls (N=530)	OR (95% CI)
	N	N	
Pilot (11) at serum sample	22	10	<b>2.54 (1.11–5.79)</b>
Pilot (11) at case diagnosis	36	23	1.72 (0.98–3.02)
Pilot (11) at serum sample and at case diagnosis	21	8	<b>2.84 (1.20–6.74)</b>
Fighter pilot (11F) at serum sample	9	3	–
Fighter pilot (11F) at case diagnosis	13	5	2.73 (0.96–7.72)
Fighter pilot (11F) at serum sample and at case diagnosis	7	2	–
Pilot/pilot trainee (11, 92T0) at serum sample	44	28	<b>1.79 (1.05–3.08)</b>
Pilot/pilot trainee (11, 92T0) at case diagnosis	46	28	<b>1.78 (1.06–3.01)</b>
Pilot/pilot trainee (11, 92T0) at serum sample and at case diagnosis	41	25	<b>1.87 (1.06–3.27)</b>
Aircraft maintenance (2A, 21A) at serum sample*	39	28	1.44 (0.86–2.39)
Aircraft maintenance (2A, 21A) at case diagnosis*	69	47	<b>1.54 (1.04–2.29)</b>
Aircraft maintenance (2A, 21A) at serum sample and at case diagnosis*	33	19	<b>1.85 (1.03–3.31)</b>

All conditional logistic regression models adjusted for number of deployments before case diagnosis (0, 1, 2+).  
 Bold indicative of statistical significance.  
 Comparison (non-exposed) group is everyone not in selected occupation/occupational group.  
 \*Models additionally adjusted for grade at case diagnosis.  
 AFSC, Air Force Specialty Code; TGCT, testicular germ cell tumour.

It is unclear what exact exposures among military pilots contribute to the elevated TGCT risk, although it has been hypothesised that taller height and higher SES may play a role.<sup>6 15 16</sup> Our sensitivity analysis adjusting for height did not significantly alter the association between pilot occupation and TGCT risk, although the sample size restricted to cases and controls without missing height information was significantly reduced. To further explore the SES hypothesis, we restricted our population to 269 cases and 262 controls with greater than high school education attainment—the association between the pilot occupation (including trainees) and TGCT risk was slightly elevated (OR=2.28, 95% CI: 1.09–4.77) compared with the original (OR=1.78, 95% CI: 1.06–3.01). It is also possible that pilots are exposed to similar chemicals as aircraft maintainers (eg, solvents, jet propellant), which may contribute to elevated TGCT risk.<sup>11 13 27</sup>

Our observed association between aircraft maintenance jobs and elevated TGCT risk is also consistent with findings from previous studies.<sup>11–13</sup> In another smaller US military study conducted among white active duty US Navy servicemen, a higher testicular cancer incidence was observed among aviation support equipment technicians compared with the total US Navy population of white active duty enlisted servicemen (standardised incidence ratio=6.9, 95% CI: 2.1–14.4); however, estimates were based on only five cases among the aviation support equipment technicians.<sup>11</sup> In a study conducted among the UK's Royal Air Force servicemen, age-adjusted incidence for testicular cancer was twofold higher among personnel working directly with aircraft (eg, engineers) than those less directly involved.<sup>12</sup> In another study, a greater than expected number of TGCT cases was observed among civilian aircraft maintenance workers; the

investigators hypothesised that exposure to an organic solvent mixture containing dimethylformamide could have contributed to the excess TGCT risk.<sup>13</sup> Aircraft maintenance workers are exposed to many different chemicals in lubricants, solvents, paint and jet propellant, and it has been hypothesised that job-related chemical exposures may contribute to TGCT development, although the biological mechanisms underlying risk are unclear.<sup>11 13 27</sup>

To our knowledge, ours is the first study to investigate the association between the firefighting and TGCT risk in the US military. We found a non-statistically significant association with elevated TGCT risk, although the sample size of firefighters was small (12 cases and 6 controls). Several previous studies of cancer risk among civilian firefighters have reported increased incidence of testicular cancer,<sup>9 10</sup> with a recent review of epidemiological studies concluding that the risk of testicular cancer was significantly higher among firefighters than the general population.<sup>10</sup> It is unclear what chemicals may contribute to the association between firefighting and testicular cancer risk, given that firefighters are exposed to a complex mixture of chemicals.<sup>9 10</sup> Recent evidence suggesting that exposure to PFAS chemicals from firefighting foams may be one contributing factor<sup>17–19</sup> should be explored further, especially in the military, given the widespread concern of drinking water contamination on the military bases where PFAS-containing firefighting foams were used.<sup>30</sup> We are currently investigating serum PFAS concentrations and TGCT risk in the same nested case–control study from which the occupational data presented here were derived.<sup>20</sup>

We found suggestions of reduced TGCT risk among servicemen in several occupational groups, including communication/electronics/wire system maintenance, aerospace medical/surgical service, intelligence and aircraft crew. Why personnel with those particular occupations may have lower risk of TGCT is not clear. It is interesting to note that aircraft crew had reduced TGCT risk, while the risk for pilots and aircraft maintainers was elevated. The explanation for this paradox could be that aircraft crew occupational group consisted of officers and enlisted personnel with jobs that have less contact time with aircraft and flightline (online supplemental table 1) than pilots and aircraft maintainers; therefore, aircrew members are likely less exposed to chemicals such as lubricants, solvents, paint and jet propellant.

Our study has several strengths. We investigated associations between a wide range of military occupations/occupational groups and TGCT risk in the largest to date TGCT case–control study among active duty USAF servicemen. The matched case–control study design allowed us to reduce potential confounding by age, race/ethnicity and timing of entering the military. We had sufficient occupational information for some of the main occupations/occupational groups of interest (ie, pilots and aircraft maintenance) at two different time points. Incident TGCT cases with confirmed pathology were ascertained from a comprehensive and centralised DoD tumour registry that also provided information on tumour morphology. To our knowledge, our study was the first that examined potential differences in the association between occupation and TGCT risk by tumour morphology.

This study also has several limitations. Because we did not have information on occupation timing and length, we made an assumption that servicemen did not change jobs in between the time points we examined and could not investigate dose–response relationships with occupational length. The strong assumption of servicemen not changing jobs between the two time periods could have resulted in exposure misclassification. This study also lacked information on exposures to specific



workplace agents. Although taller adult height is a risk factor for testicular cancer,<sup>6</sup> we were unable to adjust any of the main case-control associations for height due to the large proportion of missing information (~35%), in particular among officers (~70%). However, when we conducted analyses restricted to the subset of participants with information on height, the associations with pilots/pilot trainees, aircraft maintenance workers and fire protection workers did not change with adjustment for height, arguing against confounding as an explanation for our key findings. Because we did not have information on testicular cancer family history or relevant personal medical history (eg, cryptorchidism), we were not able to account for these risk factors in our analyses. Because we conducted multiple comparisons across many occupations, some of our results may be statistically significant due to chance, although most of our significant findings were confirmed in sensitivity analyses.

In conclusion, our findings from this large, nested case-control study of active duty USAF servicemen support the prior evidence that pilots and men with aircraft maintenance jobs have elevated risk of TGCT. We also observed evidence suggestive of an association with TGCT for servicemen with firefighting jobs. Further research in other military populations is warranted to confirm these findings, to further explore the patterns of association across quantitative metrics of occupational history and to elucidate specific occupational exposures that may contribute to the elevated TGCT risk among USAF servicemen.

**Contributors** HD-R—conceptualisation, study design, methodology, formal analysis, interpretation of data, writing (original draft). KM—interpretation of data, writing (review and editing). JR(3)—interpretation of data, writing (review and editing). CB—interpretation of data, writing (review and editing). ML—interpretation of data, writing (review and editing). PV—interpretation of data, writing (review and editing). MP—conceptualisation, investigation, writing (review and editing), interpretation of data, project administration, funding acquisition. JR(1)—conceptualisation, investigation, resources, data curation, interpretation of data, writing (review and editing), supervision, project administration, funding acquisition, guarantor.

**Funding** This research was supported by the Congressionally Directed Medical Research Program (grant #W81XWH1910444) and funding from the Intramural Research Program of the National Cancer Institute and the National Institutes of Health.

**Disclaimer** The views presented here are those of the authors and are not to be construed as official or reflecting the views of the Uniformed University of the Health Sciences, the Department of Defense, the US Air Force or the National Institutes of Health.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** All study procedures were reviewed and approved by the Uniformed Services University Institutional Review Board, the AFHSD, the DoD Joint Pathology Center, the Defense Health Agency, and the National Cancer Institute's Division of Cancer Epidemiology and Genetics.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Since these data are from military data resources, the process for obtaining assurances for public use requires authorisations by the Department of Defense. The senior author can provide additional information on the process upon request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

## ORCID iD

Hristina Denic-Roberts <http://orcid.org/0000-0001-7068-1858>

## REFERENCES

- Gurney JK, Florio AA, Znaor A, *et al.* International trends in the incidence of testicular cancer: lessons from 35 years and 41 countries. *Eur Urol* 2019;76:615–23.
- Purdue MP, Devesa SS, Sigurdson AJ, *et al.* International patterns and trends in testis cancer incidence. *Int J Cancer* 2005;115:822–7.
- Béranger R, Le Cornet C, Schüz J, *et al.* Occupational and environmental exposures associated with testicular germ cell tumours: systematic review of prenatal and life-long exposures. *PLoS One* 2013;8:e77130.
- Zhu K, Devesa SS, Wu H, *et al.* Cancer incidence in the U.S. military population: comparison with rates from the SEER program. *Cancer Epidemiol Biomarkers Prev* 2009;18:1740–5.
- Skakkebaek NE, Rajpert-De Meyts E, Main KM. Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Hum Reprod* 2001;16:972–8.
- McGlynn KA. Environmental and host factors in testicular germ cell tumors. *Cancer Invest* 2001;19:842–53.
- Oosterhuis JW, Looijenga LHJ. Human germ cell tumours from a developmental perspective. *Nat Rev Cancer* 2019;19:522–37.
- McGlynn KA, Trabert B. Adolescent and adult risk factors for testicular cancer. *Nat Rev Urol* 2012;9:339–49.
- Laroche E, L'Espérance S. Cancer incidence and mortality among firefighters: an overview of epidemiologic systematic reviews. *Int J Environ Res Public Health* 2021;18:2519.
- LeMasters GK, Genaidy AM, Succop P, *et al.* Cancer risk among firefighters: a review and meta-analysis of 32 studies. *J Occup Environ Med* 2006;48:1189–202.
- Garland FC, Gorham ED, Garland CF, *et al.* Testicular cancer in US navy personnel. *Am J Epidemiol* 1988;127:411–4.
- Foley S, Middleton S, Stitson D, *et al.* The incidence of testicular cancer in Royal Air force personnel. *Br J Urol* 1995;76:495–6.
- Ducatman AM, Conwill DE, Crawl J. Germ cell tumors of the testicle among aircraft repairmen. *J Urol* 1986;136:834–6.
- Grayson JK, Lyons TJ. Cancer incidence in United States air force aircrew, 1975–89. *Aviat Space Environ Med* 1996;67:101–4.
- Yamane GK, Johnson R. Testicular carcinoma in U.S. air force aviators: a case-control study. *Aviat Space Environ Med* 2003;74:846–50.
- Webber BJ, Tacke CD, Wolff GG, *et al.* Cancer incidence and mortality among fighter aviators in the United States air force. *J Occup Environ Med* 2022;64:71–8.
- International Agency for Research on Cancer (IARC). Some chemicals used as solvents and in polymer manufacture. In: *IARC monographs on the evaluation of carcinogenic risks to humans*. World Health Organization, 2017.
- Vieira VM, Hoffman K, Shin HM, *et al.* Perfluorooctanoic acid exposure and cancer outcomes in a contaminated community: a geographic analysis. *Environ Health Perspect* 2013;121:318–23.
- Barry V, Winquist A, Steenland K. Perfluorooctanoic acid (PFOA) exposures and incident cancers among adults living near a chemical plant. *Environ Health Perspect* 2013;121:1313–8.
- Purdue MP, Rhee J, Denic-Roberts H, *et al.* n.d. A nested case-control study of serum per- and polyfluoroalkyl substances and testicular germ cell tumors among U.S. Air Force Servicemen
- Rubertone MV, Brundage JF. The defense medical surveillance system and the Department of defense serum repository: glimpses of the future of public health surveillance. *Am J Public Health* 2002;92:1900–4.
- Perdue CL, Cost AAE, Rubertone MV, *et al.* Description and utilization of the United States department of defense serum repository: a review of published studies, 1985–2012. *PLoS One* 2015;10:e0114857.
- Air Force Personnel Center. *Air force enlisted classification directory (AFECD)*. 2021.
- Air Force Personnel Center. *Air force officer classification directory (AF OCD)*. 2021.
- Boice JD, Marano DE, Fryzek JP, *et al.* Mortality among aircraft manufacturing workers. *Occup Environ Med* 1999;56:581–97.
- Jalilian H, Ziaei M, Weiderpass E, *et al.* Cancer incidence and mortality among firefighters. *Int J Cancer* 2019;145:2639–46.
- Lipworth L, Sonderman JS, Mumma MT, *et al.* Cancer mortality among aircraft manufacturing workers: an extended follow-up. *J Occup Environ Med* 2011;53:992–1007.
- Begg CB, Zhang ZF. Statistical analysis of molecular epidemiology studies employing case-series. *Cancer Epidemiol Biomarkers Prev* 1994;3:173–5.
- Robbins AS, Pathak SR, Webber BJ, *et al.* Malignancy in U.S. air force fighter pilots and other officers, 1986–2017: a retrospective cohort study. *PLoS One* 2020;15:e0239437.
- Hu XC, Andrews DQ, Lindstrom AB, *et al.* Detection of poly- and perfluoroalkyl substances (PFAS) in U.S. drinking water linked to industrial sites, military fire training areas, and wastewater treatment plants. *Environ Sci Technol Lett* 2016;3:344–50.