



OPEN ACCESS

Original research

# Occupational exposure to respirable crystalline silica and incident idiopathic interstitial pneumonias and pulmonary sarcoidosis: a national prospective follow-up study

Inge Brosbøl Iversen <sup>1</sup>, Jesper Medom Vestergaard,<sup>1</sup> Johan Ohlander <sup>2</sup>, Susan Peters <sup>2</sup>, Elisabeth Bendstrup,<sup>3</sup> Jens Peter Ellekilde Bonde <sup>4</sup>, Vivi Schlünssen,<sup>5</sup> Jakob Hjort Bønløkke <sup>6</sup>, Finn Rasmussen,<sup>7</sup> Zara Ann Stokholm,<sup>1</sup> Michael Brun Andersen,<sup>8,9</sup> Hans Kromhout <sup>2</sup>, Henrik Albert Kolstad<sup>1,10</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/oemed-2023-108964>).

For numbered affiliations see end of article.

## Correspondence to

Dr Inge Brosbøl Iversen, Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, Aarhus, Denmark; [iniver@rm.dk](mailto:iniver@rm.dk)

Received 21 April 2023  
Accepted 26 April 2024  
Published Online First  
20 June 2024



► <https://doi.org/10.1136/oemed-2024-109599>



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Iversen IB, Vestergaard JM, Ohlander J, et al. *Occup Environ Med* 2024;**81**:279–286.

## ABSTRACT

**Background** Respirable crystalline silica is a well-known cause of silicosis but may also be associated with other types of interstitial lung disease. We examined the associations between occupational exposure to respirable crystalline silica and the risk of idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis.

**Methods** The total Danish working population was followed 1977–2015. Annual individual exposure to respirable crystalline silica was estimated using a quantitative job exposure matrix. Cases were identified in the Danish National Patient Register. We conducted adjusted analyses of exposure–response relations between cumulative silica exposure and other exposure metrics and idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis.

**Results** Mean cumulative exposure was 125  $\mu\text{g}/\text{m}^3$ -years among exposed workers. We observed increasing incidence rate ratios with increasing cumulative silica exposure for idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis. For idiopathic interstitial pneumonias and pulmonary sarcoidosis, trends per 50  $\mu\text{g}/\text{m}^3$ -years were 1.03 (95% CI 1.02 to 1.03) and 1.06 (95% CI 1.04 to 1.07), respectively. For silicosis, we observed the well-known exposure–response relation with a trend per 50  $\mu\text{g}/\text{m}^3$ -years of 1.20 (95% CI 1.17 to 1.23).

**Conclusion** This study suggests that silica inhalation may be related to pulmonary sarcoidosis and idiopathic interstitial pneumonias, though these findings may to some extent be explained by diagnostic misclassification. The observed exposure–response relations for silicosis at lower cumulative exposure levels than previously reported need to be corroborated in analyses that address the limitations of this study.

## INTRODUCTION

Interstitial lung diseases (ILDs) comprise a heterogeneous group of diseases characterised by inflammation or fibrosis of the pulmonary interstitium.<sup>1</sup> Some ILDs have known causes, among them pneumoconioses caused by inorganic dusts; hypersensitivity pneumonitis caused mainly by organic dusts;

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Respirable crystalline silica is a well-known cause of silicosis and associations with other types of interstitial lung disease have been suggested.

## WHAT THIS STUDY ADDS

⇒ This study indicates that respirable crystalline silica may be associated with more types of interstitial lung disease than silicosis, including pulmonary sarcoidosis and idiopathic interstitial pneumonias.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Our findings emphasise the need for further preventive initiatives for the millions of workers worldwide exposed to respirable crystalline silica.

autoimmune-related ILDs and ILDs caused by drugs and radiation.<sup>1</sup> Idiopathic interstitial pneumonias, on the other hand, are per definition without any known cause, even though associations with several occupational dust exposures, smoking, viruses and drugs have been reported,<sup>2–6</sup> which challenges the relevance of a separate category of idiopathic ILDs.<sup>6</sup> Sarcoidosis is a granulomatous ILD which is also considered idiopathic, but recent evidence suggests associations with occupational exposures.<sup>7,8</sup> Clinical and radiological findings of ILDs with known causes and idiopathic ILDs are not pathognomonic, may overlap and in some cases be indistinguishable.<sup>9,10</sup>

Respirable crystalline silica has been a well-known cause of ILDs for centuries, and the risk of silicosis increases with increasing cumulative exposure to respirable crystalline silica.<sup>11,12</sup> Idiopathic interstitial pneumonias and sarcoidosis have also been associated with silica exposure.<sup>2,8,13–15</sup> The health implications of respirable crystalline silica exposure affect millions of workers worldwide in construction, agriculture, mining and metal manufacturing,

where workers come into direct contact with soil and sand or process silica-containing products such as stone, concrete and glass.<sup>16 17</sup>

In this study, we examined the association between occupational exposure to respirable crystalline silica and idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis. We hypothesised that respirable crystalline silica exposure is associated with these types of ILD in an exposure-dependent manner.

## METHODS

### Study population

The study population is based on the Danish Occupational Cohort (DOC\*X), which includes all residents of Denmark with at least 1 month of gainful employment from 1976 to 2015.<sup>18</sup> For years since 1976, DOC\*X contains harmonised annual information on occupation, coded according to the 1988 International Standard Classification of Occupations (ISCO-88). We excluded workers born before 1900 or without at least 1 year of gainful employment. Within the total study population, we defined an inception population of all workers born 1956 or later with complete work histories since age 20 and a population restricted to blue-collar workers (ISCO-88 major categories 6–9) at baseline.

### ILD subtypes

We identified incident cases of idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis in the Danish National Patient Register<sup>19</sup> which contains information on all inpatient contacts with Danish hospitals since 1977 and all outpatient contacts since 1995. Diagnoses were coded according to the 8th (1977–1993) and 10th (1994–2015) versions of the International Classification of Diseases (ICD-8 and ICD-10). ICD-8 codes were only available for silicosis.

We defined idiopathic interstitial pneumonias by ICD-10 code J84; pulmonary sarcoidosis by ICD-10 codes D86.0 and D86.2; and silicosis by ICD-8 code 515.0 and ICD-10 code J62. We included only pulmonary sarcoidosis as we expected crystalline silica to be a stronger risk factor for pulmonary sarcoidosis than other types of sarcoidosis.<sup>20</sup> The J84 code is labelled 'Other interstitial pulmonary diseases' in the ICD-10 classification, but it is standard national practice in Denmark to use this code mainly for the idiopathic interstitial pneumonias.

Idiopathic pulmonary fibrosis (IPF) is the most common of the idiopathic interstitial pneumonias and has been the focus of many previous risk factor studies. Therefore, we did a subanalysis of IPF identified by ICD-10 code J84.1A.

### Respirable crystalline silica exposure assessment

Exposure assessment was based on each individuals' work history as registered in DOC\*X. When an ISCO-88 code was unknown for a certain year (13% of all employment years), we assigned the latest valid ISCO-88 code up to 5 years ago. We estimated individual quantitative respirable crystalline silica exposure for each year of employment using an ISCO-88 version of the job exposure matrix (JEM) SYN-JEM.<sup>21 22</sup> The original version of SYN-JEM contained annual exposure estimates for respirable crystalline silica for the more detailed ISCO-68 job codes by region/country, based on 23 640 personal exposure measurements from Europe and Canada. Recently, part of the exposure database was recoded to ISCO-88 codes and remodelled to create an ISCO-88 version of SYN-JEM. The ISCO-88 version of SYN-JEM is based on 15 204 personal exposure measurements

as German measurements from the MEGA-database were not available.

We used the exposure estimates for Northern European countries to create the following exposure metrics for each worker: (1) cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-year}$ ) as the sum of the annual exposure intensities; (2) highest attained exposure intensity ( $\mu\text{g}/\text{m}^3$ ) and (3) duration of exposure (years).

### Statistical analyses

We started follow-up the year following the first year of employment, as no information on month or day of employment was available. For this reason, we also lagged all independent variables by 1 year. As the National Patient Register contains information on health outcomes dating back to 1977, we started follow-up for silicosis this year. For idiopathic interstitial pneumonias and pulmonary sarcoidosis, we started follow-up in 1994 when ICD-10 was introduced in Denmark. Each worker was eligible for inclusion in analyses of all three types of ILD, so for each analysis, we followed workers until the year of first relevant diagnosis (idiopathic interstitial pneumonias, pulmonary sarcoidosis or silicosis), death, emigration, disappearance or end of follow-up on 31 December 2015, whichever came first. Workers diagnosed with each disease before begin of follow-up were excluded from the respective analysis.

We analysed associations between respirable crystalline silica and the different types of ILD separately. We used a discrete time hazard model with person-years as a unit of analysis using logistic regression, yielding incidence rate ratios (IRRs) with 95% CIs.<sup>23</sup> For each exposure metric, person-years were first categorised as either exposed or non-exposed and the exposed group was then divided into tertiles based on the distribution of the exposed person-years. Exposure tertiles were based on the study population with begin of follow-up in 1977. Analyses were performed both for the total study population (with the non-exposed group as reference) and restricted to the exposed population (with the lowest exposed tertile as reference). Analyses were adjusted for age, sex, calendar year of follow-up, education, JEM estimates of smoking, previously diagnosed connective tissue disease, medication and previously diagnosed cancer. More information on the covariates is available in online supplemental material (appendix 1). For some analyses we reduced the number of covariates, as specified in the tables, to make the models fit. All variables were treated as time-varying accounting for the change in status and increase in cumulative exposure over time.

We fitted restricted cubic splines with 95% CIs for cumulative respirable crystalline silica as a continuous variable, placing the knots at the 5, 50 and 95 percentiles.<sup>24</sup> The most exposed 2.5% were omitted to focus on the most relevant results.

Exposure tertiles were recalculated for the inception population. Due to the smaller population size, it was not possible to repeat analyses restricted to the exposed group.

We used Stata V.17 (StataCorp) for all analyses.

The study was registered at the repository of the Central Denmark Region (j.no.: 1-16-02-196-17).

## RESULTS

The total study population with begin of follow-up in 1977 comprised 5 439 728 workers with a total follow-up time of 141 508 223 person-years, while the population with begin of follow-up in 1994 comprised 5 100 706 workers with a total follow-up time of 88 145 471 person-years. Details of the establishment of the study population can be found in online supplemental figure S1.

We identified 17 869 cases of idiopathic interstitial pneumonias, 6670 cases of pulmonary sarcoidosis and 573 cases of silicosis. This corresponded to crude incidence rates per 100,000 person-years of 20.3 for idiopathic interstitial pneumonias, 7.6 for pulmonary sarcoidosis and 0.4 for silicosis.

10% of all person-years in the total study population were exposed to respirable crystalline silica (table 1). The most common exposed occupations were related to construction and farming which accounted for more than a third of all person-years contributed by the ever-exposed workers. The mean cumulative exposure was 125  $\mu\text{g}/\text{m}^3$ -years and the mean duration of exposure 8.2 years among the exposed workers. High levels of cumulative silica exposure were associated with higher age, male sex, vocational or high secondary education and previous connective tissue disease and cancer.

In the total study population, we observed increasing IRRs for idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis with increasing cumulative respirable silica exposure with trends per 50  $\mu\text{g}/\text{m}^3$ -years of 1.03 (95% CI 1.02 to 1.03), 1.06 (95% CI 1.04 to 1.07) and 1.20 (95% CI 1.17 to 1.23), respectively (table 2). Similar trends were seen when excluding the non-exposed groups. IRRs for the highest exposure group (the highest tertile) compared with the non-exposed group were 1.14 (95% CI 1.08 to 1.21), 1.61 (95% CI 1.45 to 1.78) and 2.95 (95% CI 2.31 to 3.76) for idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis. The risk of idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis increased with increasing duration of exposure. Risk patterns were less clear for the highest attained exposure.

Estimates only adjusted for age and sex are shown in online supplemental table S1.

Spline analyses showed monotonic increase in risk of idiopathic interstitial pneumonias and silicosis with increasing cumulative respirable crystalline silica exposure while the risks of pulmonary sarcoidosis increased up to around 200  $\mu\text{g}/\text{m}^3$ -years where it levelled off at about a doubled risk (figure 1).

In the inception population, we observed increasing risks with increasing cumulative respirable silica exposure for idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis (IRR 1.05 (95% CI 1.02 to 1.08), 1.07 (95% CI 1.05 to 1.10) and 1.26 (95% CI 1.17 to 1.36) per 50  $\mu\text{g}/\text{m}^3$ -years, respectively) (online supplemental table S2). The risk of silicosis also increased with increasing highest attained exposure while the risk of idiopathic interstitial pneumonias and pulmonary sarcoidosis increased with increasing duration of exposure.

In analyses restricted to blue-collar workers (online supplemental table S3), we observed patterns largely similar to those for the entire study population for pulmonary sarcoidosis and silicosis. For idiopathic interstitial pneumonias, no increasing risk with increasing exposure was observed.

In a subanalysis of IPF, we observed an IRR of 1.01 (95% CI 0.97 to 1.06) per 50  $\mu\text{g}/\text{m}^3$ -years, but the IRRs did not increase consistently with increasing exposure tertile (online supplemental table S4).

More than 80% of all exposed person-years in the total study population were accrued in occupations belonging to agriculture and construction (online supplemental table S5). Less than 1% of exposed person-years was related to work in quarries and stone-processing.

## DISCUSSION

In this study, we observed increasing risks of pulmonary sarcoidosis, idiopathic interstitial pneumonias and silicosis with increasing cumulative respirable crystalline silica exposure.

The consistent association observed between crystalline silica exposure and pulmonary sarcoidosis agrees with several previous studies.<sup>7 8 13 14</sup> Graff *et al*<sup>13</sup> observed an exposure-response relation between exposure duration and sarcoidosis. In contrast to earlier studies, however, our outcome is limited to pulmonary sarcoidosis.

An association for idiopathic interstitial pneumonias was observed in analyses of the total and inception population. Previously, a doubled risk following silica exposure was reported in a meta-analysis of four studies by Taskar *et al*<sup>2</sup> and a fivefold increase in a study by Kim *et al*.<sup>15</sup> Weaker associations have also been reported,<sup>3 4</sup> as well as no association.<sup>5</sup> None of these studies quantified silica exposure, and all studies, except Kim *et al*, investigated the association for IPF only and not idiopathic interstitial pneumonias in general. In a subanalysis, we did not find an increased risk of IPF.

Previous studies have established an exposure-response relation between cumulative respirable crystalline silica exposure and silicosis, but these studies have predominantly focused on a few occupations with high exposure levels, such as miners and diatomaceous earth workers.<sup>12 25</sup> In this study, we observed increased risk of silicosis at exposure levels significantly lower than those typically reported in prior studies, and also well below the cumulative exposure of 1000  $\mu\text{g}/\text{m}^3$ -years which would be acquired during 10 years of work at the current European Union workplace exposure limit of 100  $\mu\text{g}/\text{m}^3$ . However, a study of a cohort of industrial sand workers found a 22% increased mortality of silicosis and unspecified pneumoconiosis (as one common outcome category) for cumulative exposure levels >100 to 510  $\mu\text{g}/\text{m}^3$ -years compared with >0 to 100  $\mu\text{g}/\text{m}^3$ -years,<sup>26</sup> thus corroborating our findings at low exposure levels. Nevertheless, our cohort differs considerably from previous cohorts, as the majority of exposed person-years were accrued in the construction and farming industries where the risk of silicosis has been studied to a very limited extent.<sup>27 28</sup>

The mechanism by which silica induces pulmonary fibrosis is not fully understood.<sup>29 30</sup> When inhaled, crystalline silica particles reach the alveoli where they are phagocytosed by macrophages, thereby inducing secretion of proinflammatory substances. This, in turn, stimulates a profibrotic response with fibroblast activation, collagen secretion and subsequent fibrosis.<sup>29 30</sup> Our findings suggest that silica inhalation leads to lung injury that may be related to more clinical and radiological findings than those traditionally attributed to silicosis.

## Strengths and limitations

We studied the total Danish working population, which allowed us to study these relatively rare diseases. Some patients with minor symptoms may not be referred to hospitals, but we do not expect the likelihood of referral to be associated with exposure status as healthcare in Denmark is universal with no user payment. All diagnoses were retrieved from the National Patient Register, which is used extensively for epidemiological research. We do not know the diagnostic process leading to each diagnosis, but in Denmark, ILDs are diagnosed by specialists in pulmonology at four national ILD centres, meaning that the quality of the register data is likely high. Furthermore, a high validity of diagnoses of other pulmonary diseases,<sup>31</sup> as well as a wide

**Table 1** Distribution of person-years at risk (%) by time-varying worker characteristics and cumulative respirable crystalline silica level among 5 439 728 workers, Denmark, 1977–2015

	Cumulative respirable crystalline silica ( $\mu\text{g}/\text{m}^3\text{-years}$ )			
	0	<43.6	43.6–153.1	>153.2
	127 412 176	4 676 798	4 675 378	4 743 871
Worker characteristics	Person-years	Person-years	Person-years	Person-years
Sex				
Male	48.6	83.3	90.4	96.6
Female	51.4	16.7	9.6	3.4
Occupation*				
Armed forces	0.7	0.5	0.3	0.1
White-collar workers	46.8	16.2	9.5	7.3
Skilled blue-collar workers	8.2	17.7	18.2	34.6
Unskilled blue-collar workers	13.5	42.8	44.7	25.0
Unemployed or retired	23.3	21.2	24.6	25.5
Missing	7.5	1.5	2.7	7.4
Age				
<20	5.7	3.2	1.0	0.0
20–24	9.9	9.9	4.5	1.6
25–29	10.2	12.3	7.4	3.8
30–34	10.4	12.3	9.5	5.9
35–39	10.2	11.9	10.7	8.0
40–44	9.7	10.9	11.0	9.9
45–49	9.0	9.7	10.3	11.2
50–54	8.2	8.2	9.3	11.4
55–59	7.4	7.0	8.6	11.1
60–64	6.3	5.6	8.0	10.6
65–69	5.1	4.1	6.9	9.3
70–74	3.6	2.6	5.4	7.3
75–79	2.3	1.4	3.8	5.1
$\geq 80$	2.0	1.1	3.5	4.6
Calendar year				
1979–84	15.7	8.4	13.1	3.8
1985–94	25.6	14.9	31.0	26.6
1995–2004	27.5	28.0	25.7	35.1
2005–15	31.2	48.7	30.3	34.4
Probability of smoking				
5%–30%	34.3	35.8	19.9	18.3
31%–50%	32.2	39.6	32.3	57.1
51%–74%	33.5	24.6	47.8	24.6
Education†				
Unknown	7.7	5.7	11.6	5.5
Lower secondary	31.7	47.8	51.5	39.0
Vocational or high secondary	39.3	37.8	32.2	51.7
Short cycle higher	3.3	2.4	1.8	1.6
Medium cycle higher	12.7	4.5	2.3	1.9
Long cycle higher	5.2	1.8	0.6	0.3
Connective tissue disease				
No	99.0	99.3	99.1	98.8
Yes	1.0	0.7	0.9	1.2
Cancer				
No	97.1	97.8	97.3	96.2
Yes	2.9	2.2	2.7	3.8
Medication				
No	97.3	97.9	98.4	97.8
Yes	2.7	2.1	1.6	2.2
Duration of exposure (years)				
0	100.0	0.0	0.0	0.0

continued

Table 1 continued

	Cumulative respirable crystalline silica ( $\mu\text{g}/\text{m}^3\text{-years}$ )			
1–2	0.0	66.1	13.6	0.0
3–8	0.0	32.8	63.8	22.7
9–39	0.0	1.2	22.6	77.3

\*Grouped according to International Standard Classification of Occupations (ISCO), 1988 revision: Armed forces (ISCO-88 codes 0110), white-collar workers (ISCO-88 codes 1000–5999), skilled blue-collar workers (ISCO-88 codes 6000–7999), unskilled blue-collar workers (ISCO-88 codes 8000–9999), others (unemployed or retired).  
†Highest attained educational level.

range of other diseases,<sup>32–34</sup> has previously been observed, though the validity of ILD diagnoses has not been studied. We chose only to define pulmonary sarcoidosis by ICD-10 codes with specific mention of pulmonary involvement. Not all cases of pulmonary sarcoidosis are registered using these specific codes, meaning that we may have missed some cases, but those that are included will have pulmonary involvement with a high degree of certainty.

Diagnostic misclassification between ILD subtypes may have affected our results, as there are radiological and histological similarities between silicosis and other types of ILD, such as idiopathic interstitial pneumonias and sarcoidosis.<sup>35</sup> A detailed history of occupational exposures is part of the diagnostic programme at the Danish ILD centres, but previous silica exposure could potentially be missed in some cases, leading to actual cases of silicosis being classified as other types of ILD. Thus, diagnostic misclassification may to some extent explain the increased risks observed for idiopathic interstitial pneumonias and sarcoidosis in this study.

ILDs are a relatively rare group of diseases and age at onset is often high. Therefore, in order to identify a sufficient number of cases for all types of ILD, we included persons in our study population who had worked prior to our earliest information on occupation. Workers who held a job at baseline may constitute a healthier survivor population compared with all people who worked prior to follow-up, which may lead to bias towards the null.<sup>36</sup> To account for this issue, we repeated our analyses in the inception population born in 1956 or later with a complete work history since age 20. Even though the lower number of cases led to greater statistical uncertainty of the estimates, the overall risk patterns were largely unchanged, supporting the validity of the findings of the primary analyses.

We estimated individual exposure levels using a quantitative JEM which allowed us to perform exposure–response analyses crucial for investigation of causality. A JEM-assigned exposure is subject to non-differential misclassification because the JEM does not allow for variation in exposure within an occupation, but it will mainly lead to Berkson-type

**Table 2** Incidence rate ratios (IRR) of idiopathic interstitial pneumonias (IIPs), pulmonary sarcoidosis and silicosis following exposure to respirable crystalline silica (RCS) as assessed by a job exposure matrix, Denmark

Exposure to RCS	IIPs*		Pulmonary sarcoidosis*		Silicosis†	
	Cases	IRR (95% CI)‡	Cases	IRR (95% CI)‡	Cases	IRR (95% CI)‡
Cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-years}$ )						
0	14972	1	5527	1	369	1
<43.6	612	1.04 (0.96 to 1.13)	397	1.30 (1.17 to 1.44)	29	1.45 (0.99 to 2.13)
43.6–153.1	831	1.15 (1.07 to 1.23)	332	1.58 (1.41 to 1.76)	79	1.86 (1.45 to 2.39)
153.2–1300	1454	1.14 (1.08 to 1.21)	414	1.61 (1.45 to 1.78)	96	2.95 (2.31 to 3.76)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , including non-exposed		1.03 (1.02 to 1.03)		1.06 (1.04 to 1.07)		1.20 (1.17 to 1.23)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , excluding non-exposed§		1.03 (1.01 to 1.04)		1.03 (1.01 to 1.05)		1.16 (1.12 to 1.20)
Highest attained exposure ( $\mu\text{g}/\text{m}^3$ )¶						
0	14972	1	5527	1	369	1
<13.8	726	1.11 (1.03 to 1.20)	486	1.34 (1.22 to 1.47)	27	2.31 (1.55 to 3.45)
13.8–22.4	491	1.13 (1.03 to 1.23)	199	1.65 (1.43 to 1.90)	40	2.45 (1.76 to 3.40)
22.5–105.4	1680	1.12 (1.06 to 1.18)	458	1.57 (1.42 to 1.74)	137	2.34 (1.91 to 2.86)
Per 5 $\mu\text{g}/\text{m}^3$ , including non-exposed		1.03 (1.02 to 1.03)		1.05 (1.04 to 1.07)		1.18 (1.16 to 1.21)
Per 5 $\mu\text{g}/\text{m}^3$ , excluding non-exposed§		1.04 (1.02 to 1.05)		1.02 (1.00 to 1.04)		1.23 (1.19 to 1.27)
Duration (years)¶						
0	14972	1	5527	1	369	1
1–2	428	1.06 (0.96 to 1.17)	241	1.23 (1.08 to 1.41)	43	2.24 (1.63 to 3.07)
3–8	929	1.15 (1.08 to 1.23)	383	1.43 (1.29 to 1.59)	89	2.29 (1.81 to 2.90)
9–39	1540	1.12 (1.06 to 1.18)	519	1.67 (1.52 to 1.83)	72	2.54 (1.94 to 3.34)
Per 5 years, including non-exposed		1.03 (1.01 to 1.05)		1.17 (1.14 to 1.20)		1.33 (1.22 to 1.44)
Per 5 years, excluding non-exposed§		1.00 (0.97 to 1.03)		1.10 (1.05 to 1.14)		1.09 (0.94 to 1.25)


\*Followed from 1994 to 2015, 5 100 706 workers.

†Followed from 1977 to 2015, 5 439 728 workers.

‡Adjusted for age, sex, calendar year, education, probability of smoking, connective tissue disease, cancer and medications.

§For silicosis: not adjusted for education.

¶For silicosis: all analyses for this metric are not adjusted for education.



**Figure 1** Restricted cubic spline fits of adjusted incidence rate ratios of idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis by cumulative respirable crystalline silica. Dotted lines mark 95% CIs.

error, which will increase uncertainty of the risk estimates but not attenuate exposure–response relations.<sup>37</sup> When SYN-JEM was created, the crystalline silica exposure levels derived from the JEM were found to be of comparable magnitude to those observed in other studies.<sup>22</sup> SYN-JEM has previously confirmed known exposure–response relations for lung cancer.<sup>38</sup>

We found an increased risk of silicosis at cumulative exposure levels lower than in most existing studies. It is important to acknowledge that the cumulative exposure levels for the total study population are underestimated due to incomplete work histories. This limitation does, however, not affect the inception population. Additionally, SYN-JEM estimates geometric means rather than arithmetic means of exposure. Had we used arithmetic means, the cumulative exposure would have been approximately 60% higher. Low cumulative exposure levels in this nationwide population can also be attributed to individuals with relatively short duration of exposure, including short-term workers who are often excluded in industrial cohort studies. Nonetheless, the exposure levels within our cohort exhibit a good agreement with the above-mentioned industrial sand workers.<sup>26</sup> Finally, the use of a JEM to study diseases defined by an exposure, such as silicosis, may pose a problem, as cases of ILD are most likely to be labelled silicosis among the highest-exposed workers within an occupation, meaning that the JEM-assigned mean exposure estimates underestimate the actual exposure for diagnosed cases of silicosis. This problem does, however, not affect pulmonary sarcoidosis and idiopathic interstitial pneumonias.

Unexpectedly, we observed a number of silicosis cases in the non-exposed group. One reason for this finding may be cases who have worked in silica-exposed occupations prior to the first available register information on occupation in 1976. Furthermore, as a diagnosis of silicosis usually requires that the patient recalls previous silica exposure, there is a risk of ILD subtype misclassification if other dusts are mistaken for silica. Finally, rare silica-exposed work tasks within an otherwise unexposed occupation may not be captured in the JEM, which is a general limitation of JEM-based exposure estimation.

Using available register information, we adjusted for medications, connective tissue diseases and a proxy for radiation

therapy, which are all associated with ILD risk. We adjusted for smoking, which is a risk factor for some types of ILD,<sup>39</sup> using a JEM which has performed well in previous studies.<sup>40</sup> This means that estimates of smoking, similarly to estimates of silica exposure, are tightly connected with occupation, and therefore, adjustment for smoking may entail a risk of bias towards the null. However, risk patterns were largely unchanged when the smoking covariate was excluded from analyses. Furthermore, in an analysis of blue-collar workers expected to be more homogeneous in terms of lifestyle factors, we found risk patterns largely similar to those of the total study population, providing support that residual confounding by lifestyle factors has not significantly affected our results. We did not adjust for other occupational dust exposures associated with increased risk of ILD. Exposure to other dusts may potentially occur in most jobs assigned exposure to crystalline silica in SYN-JEM, for instance, to asbestos in construction or organic dust in agriculture,<sup>21 41</sup> and the lack of adjustment for these coexposures may have affected our results.

## CONCLUSION

In conclusion, our findings suggest that silica inhalation may be related to pulmonary sarcoidosis and idiopathic interstitial pneumonias. These findings may, however, to some extent be explained by diagnostic misclassification of silicosis. We observed exposure–response relations for silicosis at lower cumulative exposure levels than previously reported, but these findings need to be corroborated in analyses that address the limitations of this study.

## Author affiliations

- <sup>1</sup>Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, Aarhus, Denmark
- <sup>2</sup>Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands
- <sup>3</sup>Center for Rare Lung Diseases, Department of Respiratory Diseases and Allergy, Aarhus University Hospital, Aarhus, Denmark
- <sup>4</sup>Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark
- <sup>5</sup>Department of Public Health, Research Unit for Environment, Occupation and Health, Danish Ramazzini Centre, Aarhus University, Aarhus, Denmark
- <sup>6</sup>Department of Occupational and Environmental Medicine, Danish Ramazzini Centre, Aalborg University Hospital, Aalborg, Denmark
- <sup>7</sup>Department of Radiology, Aarhus University Hospital, Aarhus, Denmark
- <sup>8</sup>Department of Radiology, Copenhagen University Hospital Herlev and Gentofte, Herlev, Denmark
- <sup>9</sup>Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

<sup>10</sup>Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

**Contributors** IBI and HAK conceived and designed the study. JO, SP and HK developed the SYN-JEM ISCO-88 version. IBI and JMV established the dataset for analysis. IBI analysed the data and drafted the manuscript. All authors have contributed to interpreting the results, reviewed the paper for important intellectual content, approved the final version of the manuscript and take responsibility for the integrity of the work as a whole. IBI and HAK are guarantors of the study.

**Funding** This work was funded by grants from the Danish Working Environment Research Fund (grant no. 34-2019-09 and 47-2019-03). The original ISCO-68 SYN-JEM was developed in the SYNERGY project funded by the German Social Accident Insurance (DGUV) (grant FP 271) and was coordinated by the International Agency for Research on Cancer (IARC), the Institute for Prevention and Occupational Medicine of the DGUV, Institute of the Ruhr-University Bochum (IPA), and the Institute for Risk Assessment Sciences (IRAS) at Utrecht University. The development of the Danish Occupational Cohort (DOC\*X) was coordinated by the Department of Occupational and Environmental Medicine, Copenhagen University Hospital-Bispebjerg and Frederiksberg and funded by the Danish Working Environment Research Fund (grant no. 43-2014-03/20140016763).

**Competing interests** EB has received payment for lectures from Daiichi-Sankyo, Boehringer Ingelheim, AstraZeneca and Hoffmann-la-Roche and support for attending meetings from Boehringer Ingelheim. VS has been Chair of the Danish Quality Committee for Occupational Exposure Limits of the Danish Working Environment Authority from 2016 to 30 June 2022. MBA has received grants from the Danish Center for Lung Cancer Research, Innovation Fund Denmark and AI Signature funds from the Danish government and has received payment for lectures from Boehringer Ingelheim. HK has received a grant from Industrial Minerals Association Europe for managing the IMA-DUST Monitoring Programme. HK and SP are Editorial Board members of Occupational and Environmental Medicine. All other authors have nothing to disclose.

**Patient consent for publication** Not applicable.

**Ethics approval** No ethics approval was required. Register studies in Denmark without biological materials do not need approval from the National Committee of Health Research Ethics.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available. The data underlying this study were provided by Statistics Denmark by permission and are not publicly available. Please contact the corresponding author for any questions.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Inge Brosbøl Iversen <http://orcid.org/0009-0008-0064-4822>

Johan Ohlander <http://orcid.org/0000-0003-4279-2563>

Susan Peters <http://orcid.org/0000-0001-5662-1971>

Jens Peter Ellekilde Bonde <http://orcid.org/0000-0002-8181-3673>

Jakob Hjort Bønløkke <http://orcid.org/0000-0002-8247-6576>

Hans Kromhout <http://orcid.org/0000-0002-4233-1890>

#### REFERENCES

- 1 Wijsenbeek M, Suzuki A, Maher TM. Interstitial lung diseases. *Lancet* 2022;400:769–86.
- 2 Taskar VS, Coultas DB. Is idiopathic pulmonary fibrosis an environmental disease *Proc Am Thorac Soc* 2006;3:293–8.
- 3 Koo J-W, Myong J-P, Yoon H-K, et al. Occupational exposure and idiopathic pulmonary fibrosis: a Multicentre case-control study in Korea. *Int J Tuberc Lung Dis* 2017;21:107–12.
- 4 Awadalla NJ, Hegazy A, Elmetwally RA, et al. Occupational and environmental risk factors for idiopathic pulmonary fibrosis in Egypt: a multicenter case-control study. *Int J Occup Environ Med* 2012;3:107–16.
- 5 Abramson MJ, Murambadoro T, Alif SM, et al. Occupational and environmental risk factors for idiopathic pulmonary fibrosis in Australia: case-control study. *Thorax* 2020;75:864–9.
- 6 Wolters PJ, Blackwell TS, Eickelberg O, et al. Time for a change: is idiopathic pulmonary fibrosis still idiopathic and only Fibrotic. *Lancet Respir Med* 2018;6:154–60.
- 7 Beijer E, Meek B, Bossuyt X, et al. Immunoreactivity to metal and silica Associates with Sarcoidosis in Dutch patients. *Respir Res* 2020;21:141.
- 8 Vihlborg P, Bryngelsson I-L, Andersson L, et al. Risk of Sarcoidosis and Seropositive rheumatoid arthritis from occupational silica exposure in Swedish iron foundries: a retrospective cohort study. *BMJ Open* 2017;7:e016839.
- 9 Travis WD, Costabel U, Hansell DM, et al. An official American Thoracic society/ European respiratory society statement: update of the International Multidisciplinary classification of the idiopathic interstitial Pneumonias. *Am J Respir Crit Care Med* 2013;188:733–48.
- 10 Bendstrup E, Maher TM, Manali ED, et al. Challenges in the classification of Fibrotic ILD. *Sarcoidosis Vasc Diffuse Lung Dis* 2015;32 Suppl 1:4–9.
- 11 't Mannetje A, Steenland K, Attfield M, et al. Exposure-response analysis and risk assessment for silica and Silicosis mortality in a pooled analysis of six cohorts. *Occup Environ Med* 2002;59:723–8.
- 12 Park R, Rice F, Stayner L, et al. Exposure to crystalline silica, Silicosis, and lung disease other than cancer in Diatomaceous earth industry workers: a quantitative risk assessment. *Occup Environ Med* 2002;59:36–43.
- 13 Graff P, Larsson J, Bryngelsson I-L, et al. Sarcoidosis and silica dust exposure among men in Sweden: a case-control study. *BMJ Open* 2020;10:e038926.
- 14 Jonsson E, Järholm B, Andersson M. Silica dust and Sarcoidosis in Swedish construction workers. *Occup Med (Lond)* 2019;69:482–6.
- 15 Kim SY, Kang DM, Lee HK, et al. Occupational and environmental risk factors for chronic Fibrosing idiopathic interstitial pneumonia in South Korea. *J Occup Environ Med* 2017;59:e221–6.
- 16 Cherrie J, W, Gorman M, Seal A, et al. Shecan report, IOM: health, socio-economic and environmental aspects of possible amendments to the EU directives on the protection of workers from the risks related to the exposure to Carcinogens and Mutagens at work: Respirable Crystalline silica. 2011.
- 17 IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Arsenic, metals, fibres, and dusts. *IARC Monogr Eval Carcinog Risks Hum* 2012;100:11–465.
- 18 Flachs EM, Petersen SEB, Kolstad HA, et al. Cohort profile: DOC\*X: a nationwide Danish occupational cohort with exPosure data - an open research resource. *Int J Epidemiol* 2019;48:1413–1413k.
- 19 Lyng E, Sandegaard JL, Rebolj M. The Danish national patient register. *Scand J Public Health* 2011;39:30–3.
- 20 Ronsmans S, De Ridder J, Vandebroek E, et al. Associations between occupational and environmental exposures and organ involvement in Sarcoidosis: a retrospective case-case analysis. *Respir Res* 2021;22:224.
- 21 Peters S, Vermeulen R, Portengen L, et al. SYN-JEM: A quantitative job-exposure matrix for five lung Carcinogens. *Ann Occup Hyg* 2016;60:795–811.
- 22 Peters S, Vermeulen R, Portengen L, et al. Modelling of occupational Respirable crystalline silica exposure for quantitative exposure assessment in community-based case-control studies. *J Environ Monit* 2011;13:3262–8.
- 23 Richardson DB. Discrete time hazards models for occupational and environmental cohort analyses. *Occup Environ Med* 2010;67:67–71.
- 24 Desquilbet L, Mariotti F. Dose-response analyses using restricted cubic Spline functions in public health research. *Stat Med* 2010;29:1037–57.
- 25 Hedlund U, Jonsson H, Eriksson K, et al. Exposure-response of Silicosis mortality in Swedish iron ore miners. *Ann Occup Hyg* 2008;52:3–7.
- 26 Steenland K, Sanderson W. Lung cancer among industrial sand workers exposed to crystalline silica. *Am J Epidemiol* 2001;153:695–703.
- 27 Meijer E, Tjoer Nij E, Kraus T, et al. Pneumoconiosis and emphysema in construction workers: results of HRCT and lung function findings. *Occup Environ Med* 2011;68:542–6.
- 28 Schenker MB, Pinkerton KE, Mitchell D, et al. Pneumoconiosis from agricultural dust exposure among young California Farmworkers. *Environ Health Perspect* 2009;117:988–94.
- 29 Hoy RF, Chambers DC. Silica-related diseases in the modern world. *Allergy* 2020;75:2805–17.
- 30 Adamcakova J, Mokra D. New insights into Pathomechanisms and treatment possibilities for lung Silicosis. *Int J Mol Sci* 2021;22:4162.
- 31 Thomsen RW, Lange P, Hellquist B, et al. Validity and Underrecording of diagnosis of COPD in the Danish national patient Registry. *Respir Med* 2011;105:1063–8.
- 32 Sundbøll J, Adelborg K, Munch T, et al. Positive predictive value of cardiovascular diagnoses in the Danish national patient Registry: a validation study. *BMJ Open* 2016;6:e012832.
- 33 Ibfelt EH, Sørensen J, Jensen DV, et al. Validity and completeness of rheumatoid arthritis diagnoses in the nationwide DANBIO clinical register and the Danish national patient Registry. *Clin Epidemiol* 2017;9:627–32.
- 34 Lo B, Vind I, Vester-Andersen MK, et al. Validation of ulcerative colitis and Crohn's disease and their phenotypes in the Danish national patient Registry using a population-based cohort. *Scand J Gastroenterol* 2020;55:1171–5.

- 35 Greenberg MI, Waksman J, Curtis J. Silicosis: a review. *Dis Mon* 2007;53:394–416.
- 36 Applebaum KM, Malloy EJ, Eisen EA. Left Truncation, susceptibility, and bias in occupational cohort studies. *Epidemiology* 2011;22:599–606.
- 37 Armstrong BG. Effect of measurement error on Epidemiological studies of environmental and occupational exposures. *Occup Environ Med* 1998;55:651–6.
- 38 Ge C, Peters S, Olsson A, *et al.* Respirable crystalline silica exposure, smoking, and lung cancer subtype risks. A pooled analysis of case-control studies. *Am J Respir Crit Care Med* 2020;202:412–21.
- 39 Bae W, Lee C-H, Lee J, *et al.* Impact of smoking on the development of idiopathic pulmonary fibrosis: results from a nationwide population-based cohort study. *Thorax* 2022;77:470–6.
- 40 Bonde JPE, Flachs EM, Madsen IE, *et al.* Acute myocardial infarction in relation to physical activities at work: a nationwide follow-up study based on job-exposure Matrices. *Scand J Work Environ Health* 2020;46:3863:268–77.
- 41 Basinas I, Sigsgaard T, Kromhout H, *et al.* A comprehensive review of levels and determinants of personal exposure to dust and Endotoxin in livestock farming. *J Expo Sci Environ Epidemiol* 2015;25:123–37.



**Occupational exposure to respirable crystalline silica and incident idiopathic interstitial pneumonias and pulmonary sarcoidosis: a national prospective follow-up study**

Supplementary material

**Contents supplementary material**

Appendix 1: Description of covariates

Table S1: Results of main analysis for idiopathic interstitial pneumonias, pulmonary sarcoidosis and silicosis with age- and sex-adjusted estimates and fully adjusted estimates

Table S2: Analyses in inception population

Table S3: Analyses in population restricted to blue-collar workers at baseline

Table S4: Analyses of idiopathic pulmonary fibrosis

Table S5: Distribution of exposed person-years at risk by ISCO-88 codes

Figure S1: Flowchart of the establishment of the study population

## Appendix 1

We adjusted analyses for age divided into 5-year groups (<50, 50-54, 55-59, 60-64, 65-69, 70-74, 75-79, ≥80 years for all analyses except in the inception population where age groups were redefined (<35, 35-39, 40-44, 45-49, ≥50 years), sex (male, female) and calendar year of follow-up (1977-84, 1985-94, 1995-2004, 2005-15 for the population with begin of follow-up in 1977 and 1994-2004, 2005-2015 for the population with begin of follow-up in 1994).

We adjusted for educational level divided into the following groups: lower secondary, vocational or higher secondary, short-, medium-, and long-cycle higher education and unknown. Information on educational level for each individual was obtained from Statistics Denmark.

Smoking is a risk factor for several ILDs, including IPF (1, 2). As individual information on smoking was not available, we adjusted for estimated probability of smoking for each worker (5-30%, 31-50%, 51-74%) using a smoking job exposure matrix that was developed for the DOC\*X cohort (3). It is based on extensive Danish survey information on probability of tobacco smoking and provides sex- and calendar year-specific estimates of smoking prevalence for all ISCO-88 occupations and has been shown to prospectively predict mortality and acute myocardial infarction independent of other determinants as expected (4, 5). We assigned years without employment the same smoking probability as in the latest previous year with employment. If there was no previous information on smoking probability available due to military or non-valid ISCO-88 codes that could not be linked with the smoking JEM, we assigned the same smoking habit as in the next later period where smoking habit could be assigned.

As CTDs are known risk factors for ILD, we adjusted for previously diagnosed CTD. We identified participants with previously diagnosed CTD in the National Patient Register by ICD-8 codes 446, 712, 716 and 734 and ICD-10 codes M05, M06, M08, M09 and M30-M36.

We adjusted for redeemed prescriptions obtained from the National Prescription Registry which contains information on all prescription drugs sold in Danish pharmacies since 1994 (6). We identified the date of first redemption of one of the following drugs (classified according to the Anatomical Therapeutic Chemical (ATC) classification) which have been associated with increased risk of ILD: Antineoplastic and immunomodulating agents (ATC code L), nitrofurans derivatives (ATC code J01XE), and amiodarone (C01BD01).

The National Prescription Registry does not contain information on drugs dispensed by hospital departments, which is often the case for antineoplastic drugs. As a proxy for exposure to antineoplastic drugs and radiation therapy, which is also a risk factor for ILD, we identified participants with diagnoses of cancer, except non-melanoma skin cancer, in the National Patient Register by ICD-8 codes 140-172 and 174-209; and ICD-10 codes C00-C43 and C45-C96.

## References

1. Lee CT, Feary J, Johansson KA. Environmental and occupational exposures in interstitial lung disease. *Curr Opin Pulm Med*. 2022;28(5):414-20.
2. Bae W, Lee CH, Lee J, Kim YW, Han K, Choi SM. Impact of smoking on the development of idiopathic pulmonary fibrosis: results from a nationwide population-based cohort study. *Thorax*. 2022;77(5):470-6.
3. Bondo Petersen S, Flachs EM, Prescott EIB, Tjonneland A, Osler M, Andersen I, et al. Job-exposure matrices addressing lifestyle to be applied in register-based occupational health studies. *Occupational and environmental medicine*. 2018;75(12):890-7.

4. Flachs EM, Bondo Petersen S, Kolstad HA, Schlünssen V, Svendsen SW, Hansen J, et al. Cohort Profile: DOC\*X: a nationwide Danish occupational cohort with eXposure data - an open research resource. *International journal of epidemiology*. 2019.
5. Bonde JPE, Flachs EM, Madsen IE, Petersen SB, Andersen JH, Hansen J, et al. Acute myocardial infarction in relation to physical activities at work: a nationwide follow-up study based on job-exposure matrices. *Scand J Work Environ Health*. 2020;46(3):268-77.
6. Kildemoes HW, Sørensen HT, Hallas J. The Danish National Prescription Registry. *Scand J Public Health*. 2011;39(7 Suppl):38-41.

**Supplementary table S1.** Incidence rate ratios (IRRs) of idiopathic interstitial pneumonias (IIPs), pulmonary sarcoidosis and silicosis following exposure to respirable crystalline silica as assessed by a job exposure matrix, Denmark

Exposure	IIPs <sup>a</sup>		Pulmonary sarcoidosis <sup>a</sup>		Silicosis <sup>b</sup>	
	IRR (95% CI) <sup>c</sup>	IRR (95% CI) <sup>d</sup>	IRR (95% CI) <sup>c</sup>	IRR (95% CI) <sup>d</sup>	IRR (95% CI) <sup>c</sup>	IRR (95% CI) <sup>d</sup>
Cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-years}$ )						
0	1	1	1	1	1	1
<43.6	1.12 (1.03-1.21)	1.04 (0.96-1.13)	1.39 (1.25-1.54)	1.30 (1.17-1.44)	1.53 (1.05-2.24)	1.45 (0.99-2.13)
43.6-153.1	1.22 (1.13-1.31)	1.15 (1.07-1.23)	1.59 (1.42-1.78)	1.58 (1.41-1.76)	2.54 (1.99-3.24)	1.86 (1.45-2.39)
153.2-1300	1.15 (1.09-1.22)	1.14 (1.08-1.21)	1.56 (1.40-1.73)	1.61 (1.45-1.78)	2.30 (1.83-2.89)	2.95 (2.31-3.76)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , incl. non-exposed	1.03 (1.02-1.04)	1.03 (1.02-1.03)	1.06 (1.04-1.07)	1.06 (1.04-1.07)	1.14 (1.11-1.16)	1.19 (1.16-1.22)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , excl. non-exposed <sup>f</sup>	1.02 (1.01-1.04)	1.03 (1.01-1.04)	1.01 (1.00-1.03)	1.03 (1.01-1.05)	1.10 (1.06-1.13)	1.16 (1.12-1.20)
Highest attained exposure ( $\mu\text{g}/\text{m}^3$ ) <sup>e</sup>						
0	1	1	1	1	1	1
<13.8	1.20 (1.11-1.29)	1.11 (1.03-1.20)	1.46 (1.33-1.60)	1.34 (1.22-1.47)	1.52 (1.03-2.25)	2.31 (1.55-3.45)
13.8-22.4	1.22 (1.11-1.34)	1.13 (1.03-1.23)	1.62 (1.40-1.87)	1.65 (1.43-1.90)	2.62 (1.89-3.63)	2.45 (1.76-3.40)
22.5-105.4	1.13 (1.07-1.19)	1.12 (1.06-1.18)	1.51 (1.36-1.66)	1.57 (1.42-1.74)	2.33 (1.91-2.85)	2.34 (1.91-2.86)
Per 5 $\mu\text{g}/\text{m}^3$ , incl. non-exposed	1.03 (1.03-1.04)	1.03 (1.02-1.03)	1.05 (1.04-1.06)	1.05 (1.04-1.07)	1.17 (1.15-1.20)	1.18 (1.16-1.20)
Per 5 $\mu\text{g}/\text{m}^3$ , excl. non-exposed <sup>f</sup>	1.04 (1.03-1.05)	1.04 (1.02-1.05)	1.00 (0.98-1.02)	1.02 (1.00-1.04)	1.23 (1.19-1.27)	1.23 (1.19-1.27)
Duration (years) <sup>e</sup>						
0	1	1	1	1	1	1
1 – 2	1.13 (1.02-1.24)	1.06 (0.96-1.17)	1.29 (1.13-1.47)	1.23 (1.08-1.41)	2.66 (1.94-3.65)	2.24 (1.63-3.07)
3 – 8	1.22 (1.14-1.30)	1.15 (1.08-1.23)	1.47 (1.32-1.63)	1.43 (1.29-1.59)	2.68 (2.13-3.39)	2.29 (1.81-2.90)
9 – 39	1.14 (1.08-1.20)	1.12 (1.06-1.18)	1.67 (1.52-1.83)	1.67 (1.52-1.83)	1.68 (1.30-2.17)	2.54 (1.94-3.34)
Per 5 years, incl. non-exposed	1.04 (1.02-1.06)	1.03 (1.01-1.05)	1.18 (1.15-1.21)	1.17 (1.14-1.20)	1.19 (1.11-1.28)	1.40 (1.29-1.51)
Per 5 years, excl. non-exposed <sup>f</sup>	0.98 (0.96-1.01)	1.00 (0.97-1.03)	1.10 (1.06-1.15)	1.10 (1.05-1.14)	0.87 (0.77-0.98)	1.09 (0.94-1.25)

<sup>a</sup>Followed from 1994-2015, 5,100,706 workers<sup>b</sup>Followed from 1977-2015, 5,439,728 workers<sup>c</sup>Adjusted for age and sex<sup>d</sup>Adjusted for age, sex, calendar year, education, probability of smoking, connective tissue disease, cancer and medications<sup>e</sup>For silicosis: all analyses for this metric are not adjusted for education<sup>f</sup>For silicosis: not adjusted for education

**Supplementary table S2.** Incidence rate ratios (IRR) of idiopathic interstitial pneumonias (IIPs), pulmonary sarcoidosis and silicosis following exposure to respirable crystalline silica as assessed by a job exposure matrix among workers born 1956 or later, Denmark

Exposure	IIPs <sup>a</sup>		Pulmonary sarcoidosis <sup>a</sup>		Silicosis <sup>b</sup>	
	Cases	IRR (95% CI) <sup>c</sup>	Cases	IRR (95% CI) <sup>c</sup>	Cases <sup>e</sup>	IRR (95% CI) <sup>d</sup>
Cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-years}$ )						
0	2128	1	3353	1	18	1
<24.6	177	1.19 (1.02-1.39)	312	1.34 (1.19-1.51)	n.r.	1.57 (0.36-6.93)
24.6-89.5	121	1.34 (1.11-1.62)	225	1.67 (1.45-1.91)	n.r.	1.91 (0.55-6.68)
>89.5	88	1.45 (1.16-1.80)	154	1.62 (1.37-1.91)	n.r.	4.34 (1.84-10.25)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , incl. non-exposed		1.05 (1.02-1.08)		1.07 (1.05-1.10)		1.26 (1.17-1.36)
Highest attained exposure ( $\mu\text{g}/\text{m}^3$ )						
0	2128	1	3353	1	18	1
<9.7	230	1.30 (1.13-1.49)	391	1.42 (1.27-1.58)	n.r.	1.33 (0.30-5.91)
9.7-20.0	58	1.17 (0.89-1.52)	129	1.89 (1.58-2.25)	n.r.	2.97 (1.06-8.33)
>10.0	98	1.34 (1.09-1.66)	171	1.44 (1.23-1.68)	n.r.	3.91 (1.56-9.78)
Per 5 $\mu\text{g}/\text{m}^3$ , incl. non-exposed		1.04 (1.02-1.07)		1.06 (1.04-1.08)		1.21 (1.11-1.32)
Duration (years)						
0	2128	1	3353	1	18	1
1	103	1.11 (0.91-1.35)	186	1.26 (1.08-1.46)	n.r.	1.50 (0.20-11.34)
2-5	158	1.37 (1.16-1.61)	271	1.48 (1.30-1.67)	n.r.	0.65 (0.09-4.97)
6-39	125	1.38 (1.14-1.66)	234	1.78 (1.55-2.04)	n.r.	4.62 (2.06-10.34)
Per 5 years, incl. non-exposed		1.11 (1.05-1.17)		1.20 (1.16-1.25)		1.52 (1.23-1.87)

n.r.: not reported, cells with less than five cases.

<sup>a</sup>Followed from 1994-2015, 2,955,863 workers<sup>b</sup>Followed from 1977-2015, 2,980,445 workers<sup>c</sup>Adjusted for age, sex, calendar year, education, probability of smoking, connective tissue disease, cancer and medications<sup>d</sup>Adjusted for age, sex, calendar year (1977-94, 1995-2004, 2005-15), probability of smoking, connective tissue disease and medications<sup>e</sup>Total number of silicosis cases: 32

**Supplementary table S3.** Incidence rate ratios (IRR) of idiopathic interstitial pneumonias (IIPs), pulmonary sarcoidosis and silicosis following exposure to respirable crystalline silica as assessed by a job exposure matrix among blue collar workers, Denmark

Exposure	IIPs <sup>a</sup>		Pulmonary sarcoidosis <sup>a</sup>		Silicosis <sup>a</sup>	
	Cases	IRR (95% CI) <sup>c</sup>	Cases	IRR (95% CI) <sup>c</sup>	Cases	IRR (95% CI) <sup>d</sup>
Cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-years}$ )						
0	3213	1	1501	1	256	1
<43.6	329	1.06 (0.95-1.19)	210	1.14 (0.99-1.32)	22	1.40 (0.91-2.17)
43.6-153.1	376	1.14 (1.02-1.27)	229	1.46 (1.27-1.68)	66	2.62 (2.00-3.44)
153.2-1300	1019	1.01 (0.94-1.10)	358	1.51 (1.33-1.70)	86	3.42 (2.65-4.41)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , incl. non-exposed		1.01 (1.00-1.02)		1.05 (1.03-1.06)		1.13 (1.09-1.17)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , excl. non-exposed		1.02 (1.00-1.03)		1.03 (1.01-1.05)		1.08 (1.02-1.15)
Highest attained exposure ( $\mu\text{g}/\text{m}^3$ )						
0	3213	1	1501	1	256	1
<13.8	472	1.12 (1.01-1.24)	286	1.18 (1.04-1.35)	18	1.65 (1.02-2.69)
13.8-22.4	242	1.03 (0.90-1.18)	154	1.68 (1.42-1.99)	33	1.73 (1.20-2.48)
22.5-105.4	1010	1.02 (0.95-1.10)	357	1.45 (1.29-1.64)	123	3.34 (2.69-4.15)
Per 5 $\mu\text{g}/\text{m}^3$ , incl. non-exposed		1.02 (1.01-1.02)		1.04 (1.03-1.06)		1.15 (1.13-1.17)
Per 5 $\mu\text{g}/\text{m}^3$ , excl. non-exposed		1.02 (1.00-1.04)		1.01 (0.99-1.04)		1.17 (1.12-1.21)
Duration (years)						
0	3213	1	1501	1	256	1
1 – 2	199	1.09 (0.94-1.26)	129	1.13 (0.95-1.36)	39	2.43 (1.73-3.40)
3 – 8	360	1.08 (0.97-1.21)	228	1.32 (1.14-1.52)	76	2.51 (1.94-3.25)
9 – 39	1165	1.03 (0.96-1.11)	440	1.51 (1.35-1.69)	59	2.99 (2.20-4.07)
Per 5 years, incl. non-exposed		1.00 (0.98-1.03)		1.13 (1.10-1.17)		1.40 (1.28-1.53)
Per 5 years, excl. non-exposed		0.98 (0.95-1.02)		1.08 (1.03-1.14)		1.12 (0.96-1.30)

<sup>a</sup>Followed from 1994-2015, 1,375,815 workers<sup>b</sup>Followed from 1977-2015, 1,454,098 workers<sup>c</sup>adjusted for age, sex, calendar year, education, probability of smoking, connective tissue disease, cancer and medications<sup>d</sup>Adjusted for age, sex, calendar year, probability of smoking, connective tissue disease, cancer and medications

**Supplementary table S4.** Incidence rate ratios (IRR) of idiopathic pulmonary fibrosis (IPF) following exposure to respirable crystalline silica as assessed by a job exposure matrix among 5,100,706 workers, Denmark 1994-2015

Exposure	IPF	
	Cases	IRR (95% CI) <sup>a</sup>
Cumulative exposure ( $\mu\text{g}/\text{m}^3\text{-years}$ )		
0	321	1
<43.6	13	0.91 (0.52-1.59)
43.6-153.1	20	1.35 (0.85-2.15)
153.2-1300	37	1.00 (0.70-1.44)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , incl. non-exposed		1.01 (0.97-1.06)
Per 50 $\mu\text{g}/\text{m}^3\text{-years}$ , excl. non-exposed <sup>b</sup>		1.03 (0.96-1.10)
Highest attained exposure ( $\mu\text{g}/\text{m}^3$ )		
0	321	1
<13.8	21	1.19 (0.76-1.88)
13.8-22.4	12	1.23 (0.68-2.22)
22.5-105.4	37	0.95 (0.66-1.36)
Per 5 $\mu\text{g}/\text{m}^3$ , incl. non-exposed		1.00 (0.96-1.05)
Per 5 $\mu\text{g}/\text{m}^3$ , excl. non-exposed <sup>b</sup>		0.99 (0.91-1.07)
Duration (years)		
0	321	1
1 – 2	8	0.92 (0.45-1.86)
3 – 8	19	1.15 (0.72-1.85)
9 – 39	43	1.05 (0.75-1.48)
Per 5 years, incl. non-exposed		1.02 (0.92-1.13)
Per 5 years, excl. non-exposed <sup>b</sup>		1.06 (0.89-1.27)

<sup>a</sup>Adjusted for age, sex, calendar year, education, probability of smoking, connective tissue disease, cancer and medications

<sup>b</sup>Not adjusted for education



**Table S5.** Distribution of exposed person-years at risk by ISCO-88 codes in the total study population.

ISCO-88	Description	Person-years	Percent
2452	Sculptors, painters and related artists	48,895	0.90
6110	Market gardeners and crop growers	324	0.01
6111	Field crop and vegetable growers	82,577	1.53
6112	Tree and shrub crop growers	213,010	3.93
6113	Gardeners, horticultural and nursery growers	34,610	0.64
6130	Market-oriented crop and animal producers	231,362	4.27
7100	Extraction and building trades workers	62,219	1.15
7110	Miners, shotfirers, stone cutters and carvers	215	0.00
7111	Miners and quarry workers	861	0.02
7112	Shotfirers and blasters	313	0.01
7113	Stone splitters, cutters and carvers	9,061	0.17
7120	Building frame and related trades workers	654,767	12.09
7121	Builders, traditional materials	13,933	0.26
7122	Bricklayers and stonemasons	442,314	8.17
7123	Concrete placers, concrete finishers and related workers	6,975	0.13
7129	Building frame and related trades workers not elsewhere classified	61,477	1.14
7131	Roofers	28,441	0.53
7132	Floor layers and tile setters	26,692	0.49
7133	Plasterers	5,510	0.10
7143	Building structure cleaners	31,624	0.58
7211	Metal moulders and coremakers	24,236	0.45
7321	Abrasive wheel formers, potters and related workers	14,530	0.27
8110	Mining- and mineral-processing plant operators	21	0.00
8111	Mining-plant operators	117	0.00
8112	Mineral-ore- and stone-processing-plant operators	9,835	0.18
8113	Well drillers and borers and related workers	13,032	0.24
8130	Glass, ceramics and related plant operators	3,035	0.06
8131	Glass and ceramics kiln and related machine operators	56,651	1.05
8139	Glass, ceramics and related plant operators not elsewhere classified	13,409	0.25
8210	Metal- and mineral-products machine operators	65	0.00
8212	Cement and other mineral products machine operators	47,480	0.88
8223	Metal finishing-, plating- and coating-machine operators	48,335	0.89
8330	Agricultural and other mobile-plant operators	7,264	0.13
8331	Motorised farm and forestry plant operators	42,992	0.79
8332	Earth-moving- and related plant operators	101,738	1.88
9160	Garbage collectors and related labourers	58,473	1.08
9161	Garbage collectors	28,266	0.52
9162	Sweepers and related labourers	27,572	0.51
9211	Farm-hands and labourers	1,457,115	26.91
9300	Labourers in mining, construction, manufacturing and transport	42,620	0.79
9310	Mining and construction labourers	149,577	2.76
9311	Mining and quarrying labourers	2,695	0.05
9312	Construction and maintenance labourers: roads, dams and similar constructions	617,009	11.40
9313	Building construction labourers	693,234	12.80
<b>Total</b>		<b>5,414,481</b>	<b>100.00</b>

**Supplementary figure S1.** Establishment of the study population.