

ORIGINAL RESEARCH

Association between SARS-CoV-2 infection, exposure risk and mental health among a cohort of essential retail workers in the USA

Fan-Yun Lan,^{1,2} Christian Suharlim,³ Stefanos N Kales,^{1,4} Justin Yang ^{1,5}

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2020-106774>).

¹Environmental & Occupational Medicine & Epidemiology Program, Department of Environmental Health, Harvard University T H Chan School of Public Health, Boston, Massachusetts, USA

²Department of Occupational and Environmental Medicine, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan, Taiwan

³Management Sciences for Health, Medford, Massachusetts, USA

⁴Department of Occupational Medicine, Cambridge Health Alliance, Harvard Medical School, Cambridge, Massachusetts, USA

⁵Department of General Internal Medicine, Boston University School of Medicine, Boston, Massachusetts, USA

Correspondence to

Dr Justin Yang, Environmental Health, Harvard University T H Chan School of Public Health, Boston, MA 02115, USA; justin.yang@mail.harvard.edu

Received 8 June 2020

Revised 11 September 2020

Accepted 26 September 2020



© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Lan F-Y, Suharlim C, Kales SN, *et al.* *Occup Environ Med Epub ahead of print*: [please include Day/Month/Year]. doi:10.1136/oemed-2020-106774

ABSTRACT

Objectives To investigate SARS-CoV-2 (the virus causing COVID-19) infection and exposure risks among grocery retail workers, and to investigate their mental health state during the pandemic.

Methods This cross-sectional study was conducted in May 2020 in a single grocery retail store in Massachusetts, USA. We assessed workers' personal/occupational history and perception of COVID-19 by questionnaire. The health outcomes were measured by nasopharyngeal SARS-CoV-2 reverse transcriptase PCR (RT-PCR) results, General Anxiety Disorder-7 (GAD-7) and Patient Health Questionnaire-9 (PHQ-9).

Results Among 104 workers tested, 21 (20%) had positive viral assays. Seventy-six per cent positive cases were asymptomatic. Employees with direct customer exposure had an odds of 5.1 (95% CI 1.1 to 24.8) being tested positive for SARS-CoV-2 after adjustments. As to mental health, the prevalence of anxiety and depression (ie, GAD-7 score >4 or PHQ-9 score >4) was 24% and 8%, respectively. After adjusting for potential confounders, those able to practice social distancing consistently at work had odds of 0.3 (95% CI 0.1 to 0.9) and 0.2 (95% CI 0.03 to 0.99) screening positive for anxiety and depression, respectively. Workers commuting by foot, bike or private cars were less likely to screen positive for depression (OR 0.1, 95% CI 0.02 to 0.7).

Conclusions In this single store sample, we found a considerable asymptomatic SARS-CoV-2 infection rate among grocery workers. Employees with direct customer exposure were five times more likely to test positive for SARS-CoV-2. Those able to practice social distancing consistently at work had significantly lower risk of anxiety or depression.

INTRODUCTION

WHO declared COVID-19 as a pandemic on 11 March 2020.¹ Since then, accumulating evidence has shown the transmission capability of SARS-CoV-2, the virus causing COVID-19, not just from symptomatic patients but from asymptomatic carriers.²⁻⁴ Interventions have been implemented worldwide to minimise transmission, including social distancing, travel bans, stay-at-home orders and school and non-essential business closures.^{5,6} All measures are intended to reduce contact and to prevent transmission, especially when the index patients are in subclinical stage of SARS-CoV-2 infection.⁷ While most community residents benefit from these risk reduction policies, certain essential employees, such

Key messages

What is already known about this subject?

- The health of essential workers during the COVID-19 pandemic is of great public and media interests.
- Research, however, has largely focused on healthcare workers with relatively limited literature investigating non-healthcare essential workers.
- Previous studies suggested essential workers are not able to benefit from mitigation policies.
- Their occupational exposures increase their own risk to SARS-CoV-2 infection, and increase the risk of secondary transmissions to their colleagues, families and communities.

What are the new findings?

- The present study fills in the knowledge gap of COVID-19 impacts on grocery/retail market workers during the pandemic, from both physical and psychological perspectives.
- In this single store sample (n=104), we found an alarming infection rate of 20% positive SARS-CoV-2 RT-PCR assay result among these workers and the majority (76%) of them were asymptomatic at the time of testing.
- Furthermore, employees with direct customer exposure were five times more likely to test positive for SARS-CoV-2.
- Our study also found the inability to practice social distancing consistently at work was a significant risk factor for anxiety and depression.
- At the same time, commuting to work by public transportation/shared rides was significantly associated with depressive state.

as healthcare workers (HCWs), first responders and retail workers, continue to experience potential SARS-CoV-2 exposure risk due to the nature of their job.⁸ Furthermore, once essential workers are infected with SARS-CoV-2, they may become a significant transmission source for the community they serve.⁹

The psychological stress associated with working during the COVID-19 pandemic is also of great public interest.¹⁰ Studies have indicated pandemic awareness, infection fear and family concerns

Key messages

How might this impact on policy or clinical practice in the foreseeable future?

- ▶ This is the first study to demonstrate the significant asymptomatic infection rate, exposure risks and associated psychological distress of grocery retail essential workers during the pandemic, which supports the policy recommendations that employers and government officials should take actions on implementing preventive strategies and administrative arrangements, such as methods to reduce interpersonal contact, repeat and routine SARS-CoV-2 employee testing, to ensure the health and safety of essential workers.
- ▶ Our significant mental health finding calls for action in providing comprehensive employee assistance services to help essential workers cope with the psychological distress during the COVID-19 pandemic.

contribute significantly to essential workers' mental distress during an emerging disease pandemic.^{11 12}

Pioneering COVID-19 studies on essential workers have largely focused on HCWs. Studies showed the attack rates of SARS-CoV-2 among HCWs in early outbreaks ranged from 0% to 14%, with fever and loss of smell/taste being the best predictors of the disease.^{13 14} In terms of mental health, about half of the HCWs included in one study reported anxiety and depressive symptoms with psychological stress risk factors including living in areas with higher prevalence or being frontline HCWs.¹⁵

While HCWs have been widely discussed in COVID-19-related research, there are relatively limited studies investigating other essential workers. A recent publication looking at six Asian countries showed that various non-HCWs were also affected during early COVID-19 transmission, with service and sales workers comprising 18% of possible work-related cases.⁹ While previous studies have reported SARS-CoV-2 cluster infections in supermarket settings,^{16 17} no study has examined the SARS-CoV-2 exposure risks or psychological stress among grocery retail essential employees. Therefore, we conducted this study aiming to investigate: 1) SARS-CoV-2 infection rate, transmission and exposure risks among grocery retail employees, 2) their use of personal protective equipment (PPE) and perception on COVID-19 and 3) their mental health state during the COVID-19 pandemic.

METHODS**Study design and study population**

This cross-sectional study is reported according to the Strengthening the Reporting of Observational Studies in Epidemiology guideline.¹⁸ We used secondary data from a COVID-19 testing tent site that included information collected from 104 adults employed at one grocery retail store in the greater Boston area of Massachusetts, USA as part of a city-mandated group testing. Clinical evaluation and nasopharyngeal swab sampling were conducted on each individual over three consecutive days in early May 2020. All workers older than 18 years sent by the store and presented for testing were included in this study (100% response rate).

SARS-CoV-2 RT-PCR specimen collection and testing

The specimens were collected using nasopharyngeal swab inside the designated COVID-19 testing tent. A trained physician

performed the swabbing procedure and transferred each specimen to a 3 mL vial with viral transport media. The samples were then transported to Quest Diagnostic laboratory in Marlborough, Massachusetts, where real-time, reverse-transcriptase-PCR (RT-PCR) diagnostic panels were conducted to detect SARS-CoV-2. All sampling, specimen storage, transportation and testing procedures followed the guidelines of the US Centers for Disease Control and Prevention.¹⁹

Questionnaire survey

As part of the group testing procedure, participants' basic demographic information, SARS-CoV-2-related exposure information, PPE usage and mental health surveys were collected through a paper-based questionnaire completed on site prior to testing.

The basic information section of questionnaire included age, sex, race/ethnicity and medical history including past medical problems, prescription medication history, smoking status, alcohol intake, recreational drug use history and primary care physician information. For past medical issues, participants responded to a checklist which included the following diseases: chronic obstructive pulmonary disease/emphysema, asthma, heart disease, high cholesterol, high blood pressure, diabetes, HIV, hepatitis C, cancer and other(s).

The following questions were included for employment history: most recent job position(s) at the store in the past month, full-time/part-time employment status, work hours per week (<20 hours, 20–39 hours, 40 hours and above), average length of shifts, additional employment(s) outside this retail store and transportation method(s) to work (by foot or bike, private car, public transportation, shared rides or others). Workers selected their job position(s) from the following choices: cashier, front end associate, cart attendant, janitorial crew, stocker, backroom, receiving, sales associate, fresh food associate, supervisor and/or specialised roles. Participants were given the choice to answer with free text for some other position if not listed as above. Employees were asked to identify any additional employment(s) in the following categories: healthcare, drivers and transport, services and sales, cleaning and domestic, public safety, restaurant/fast food, others.⁹

As to COVID-19-related information, participants indicated new-onset symptoms within the past 1–2 weeks as a yes or no to a checklist of 11 common COVID-19 symptoms, including fever/chills, headache, running nose, sore throat, cough (acute, new onset, dry or productive), shortness of breath, loss of taste or smell, diffuse body ache, fatigue/feeling run down, nausea, diarrhoea. If participants answered yes to any of the above symptom(s), they were asked to indicate symptom onset. Participants were asked if they had been exposed to anyone that has confirmed SARS-CoV-2 in the past 14 days. If they answered yes, they were asked of whom the exposure was (colleague, friend, family/relatives) and how many days ago the exposure occurred.

Information on mental health was recorded using two validated screening tools on depression and anxiety: Patient Health Questionnaire-9 (PHQ-9)²⁰ and General Anxiety Disorder-7 (GAD-7).²¹ For PHQ-9, a total score of no higher than 4 indicates no or minimal depression, with a total PHQ-9 score ranging from 0 to 27. The score of GAD-7 ranges from 0 to 21. A GAD-7 score of no higher than 4 indicates no or minimal anxiety. Participants were also asked to self-identify any history of depression and/or anxiety.

Social distancing, PPE usage, COVID-19 prevention knowledge score and COVID-19 pandemic perception score

Participants answered a Likert scale, from never (one) to always (five), for four questions that assessed employee's practice of social distancing and PPE use. Participants answered another Likert scale with six statements, from completely disagree (one) to completely agree (five), which captured the workers' knowledge on PPE and self-perceptions toward COVID-19 pandemic. Both employee's PPE knowledge and COVID-19 perception were then tabulated to a score ranging from 3 to 15. A complete list of questions is included in Online-Only Supplement 1.

Customer exposure categorisation

Employees' job position was classified into two categories: those with significant face-to-face, direct exposure to customers and those without significant customer exposure. Employees with direct customer exposure include cashier, front end associate, sales associate, fresh food associate, cart attendant, janitorial crew, supervisor and manager of all levels. Those without direct customer exposure include stocker, backroom, receiving and maintenance.

Study participants

The COVID-19 testing was conducted as part of a city-mandated group testing, independent to this research. The existing medical records collected for the city testing were de-identified at the primary clinical site prior to analysis. Therefore, the study of de-identified data received a non-human research determination by the Management Sciences for Health (SC#0012020).

Statistical analysis

We performed univariate analyses to compare the workers' characteristics by their SARS-CoV-2 RT-PCR testing results, anxiety and depression status. For binary variables, Pearson's χ^2 test with Yates' continuity correction was performed, while for variables with at least one cell count less than five, Fisher's exact test was conducted instead. As to continuous variables, data were examined by Q-Q plots and determined if they followed normal distribution beforehand. Then we performed parametric t-test or non-parametric Wilcoxon rank sum test, as appropriate.

Logistic regression models and models adjusting for potential confounders were further built. Due to the small sample size and event numbers, we used the inverse probability weighting (IPW) method to avoid inflated SEs of the parameter estimates.²² The IPW was calculated based on the selected variables determined from the univariate analyses results. Extreme weights (below the 5th and above the 95th percentile) were truncated as an additional sensitivity analysis. ORs with 95% CIs were presented.

We performed secondary sensitivity analysis according to employees' job titles. Employees' job position(s) were initially categorised into positions with greater direct customer exposure versus those without. In the sensitivity analysis, we categorised the jobs into supervisory positions vs non-supervisory positions.

All p values reported are two-tailed. A p value <0.05 was considered statistically significant. We used R software (V3.6.3) to conduct statistical analyses.

RESULTS

In table 1, we presented the characteristics of all tested employees stratified by SARS-CoV-2 RT-PCR assay results. Among the 104

Table 1 Characteristics of retail essential employees in a single grocery store in Massachusetts, USA by SARS-CoV-2 (the virus causing COVID-19) RT-PCR assay testing results

	Overall (N=104)	Positive SARS-CoV-2 RT-PCR assay (N=21)	Negative SARS-CoV-2 RT-PCR assay (N=83)	P value
Age, mean (SD)	49.0 (14.1)	49.2 (14.4)	49.0 (14.2)	0.954
Female, n (%)	49 (47.1%)	11 (52.4%)	38 (45.8%)	0.767
Non-Caucasian, n (%)	64 (61.5%)	14 (66.7%)	50 (60.2%)	0.283
Cigarette smoker, n (%)	25 (24.0%)	1 (4.8%)	24 (28.9%)	0.022*
Daily alcohol consumption, n (%)	8 (7.7%)	0	8 (9.6%)	0.354*
Marijuana use, n (%)	14 (13.5%)	2 (9.5%)	12 (14.5%)	0.730*
Self-reported exposure to SARS-CoV-2-positive individual(s) in the past 14 days, n (%)	24 (23.1%)	4 (19.0%)	20 (24.1%)	0.776*
Job positions with direct customer exposure at store†, n (%)	68 (65.4%)	19 (90.5%)	49 (59.0%)	0.009*
Full-time employment status, n (%)	73 (70.2%)	16 (76.2%)	57 (68.7%)	0.685
Residential area SARS-CoV-2 prevalence (per 100 000), geometric mean (geometric SD)	1106.0 (1.5)	1292.8 (1.63)	1063.2 (1.4)	0.179‡
Ability to practice social distancing consistently at work, count (%)	69 (66.3%)	13 (61.9%)	56 (67.5%)	0.930
Using gloves consistently at work, count (%)	80 (76.9%)	19 (90.5%)	61 (73.5%)	0.068*
Wearing face mask consistently at work, count (%)	95 (91.3%)	20 (95.2%)	75 (90.4%)	0.596*
Wearing face mask consistently outside of work, count (%)	81 (77.9%)	18 (85.7%)	63 (75.9%)	0.348*
Commute to work by foot, bike or private car	90 (86.5%)	19 (90.5%)	71 (85.5%)	0.730*
PPE knowledge score, median (IQR)	15 (14–15)	15 (14–15)	15 (14–15)	0.966 ‡
COVID-19 perception score, median (IQR)	12 (11–15)	13 (11–15)	12 (11–14)	0.510 ‡
GAD-7 score, median (IQR)	0 (0–4)	1 (0–4.5)	0 (0–4)	0.660‡
PHQ-9 score, median (IQR)	0 (0–2)	0 (0–1)	0 (0–2)	0.733‡
Employee has an assigned primary care provider, n (%)	77 (74.0%)	17 (81.0%)	60 (72.3%)	0.584*
Requested mental health support on survey, n (%)	14 (13.5%)	3 (14.3%)	11 (13.3%)	0.999*

*Statistics derived from Fisher's exact test.

†Direct customer exposure positions include cashier, front end associate, sales associate, fresh food associate, cart attendant, janitorial crew, supervisor and manager of all levels. These are in contrast to positions mainly dealing with consumer goods or the environment, such as stocker, backroom, receiving and maintenance.

‡Statistics derived from Wilcoxon rank sum test with continuity correction.

GAD-7, Generalised Anxiety Disorder 7-item scale; PHQ-9, Patient Health Questionnaire-9; PPE, personal protective equipment.

grocery retail employees that underwent testing and completed the survey, 47% were female with an average age of 49 years. The majority (62%) of employees in this retail store were non-Caucasian minorities. Twenty-one out of 104 employees tested positive for SARS-CoV-2 indicating a point prevalence of 20%. Among these SARS-CoV-2-positive employees, 91% of them had a job position with significant direct customer exposure compared with 59% among the SARS-CoV-2-negative employees ($p=0.009$). Seventy-six per cent of workers with positive tests were asymptomatic. Among the 25 smokers, only one tested positive for SARS-CoV-2 ($p=0.022$). We did not observe statistical difference of SARS-CoV-2 status associated with protective behaviour (social distancing, use of gloves and/or masks and avoid commuting by public transportation or shared rides), nor did we find significant differences in PPE knowledge, COVID-19 perception and mental health status between SARS-CoV-2-positive and SARS-CoV-2-negative employees.

Table 2 shows the distributions of workers' characteristics, comparing those with at least mild anxiety versus those reporting no or minimal anxiety. Ninety-nine out of 104 workers (95%) completed the GAD-7 questionnaire, with 24 workers (24%) reporting at least mild anxiety. We observed no statistical differences to anxiety by age, gender, smoking, alcohol consumption, marijuana use, possible SARS-CoV-2 exposure, job position, commuting method and PPE use. Only 46% of workers with anxiety reported they were able to practice social distancing consistently at work, whereas the majority (76%) of those without reported anxiety were able to do so at work ($p=0.009$). Employees screening positive for anxiety also reported less consistent mask use (63%) comparing with those screened negative for anxiety (84%), although this result did not reach statistical significance ($p=0.072$). The COVID-19 pandemic perception score, which mainly evaluated the extent of worries on getting oneself and one's family infected due to work, were equally high among employees who screened positive for anxiety by GAD-7 and those who did not (median score 13 vs 12, $p=0.09$).

As to depression, there were 8 out of 99 (8%) who screened positive for at least mild depression (table 3). Workers who

reported at least mild depression recorded higher proportion of possible SARS-CoV-2 exposure in the past 14 days compared with those without depression (63% vs 21%, $p=0.019$). Workers who screened positive for depression by PHQ-9 were less likely to practice social distancing consistently at work and more likely to commute by public transportation or shared rides, compared with those without depression (25% vs 73% and 50% vs 11%, $p=0.010$ and $p=0.013$, respectively).

Employees with direct customer exposure were five times more likely to test positive on SARS-CoV-2 RT-PCR assay comparing with those without direct customer exposures (OR 5.1, 95% CI 1.1 to 24.8) after adjusting for age, gender, smoking and SARS-CoV-2 community prevalence in workers' residential cities (table 4). While cigarette smokers had an 90% risk reduction in having positive SARS-CoV-2 RT-PCR assay result in the crude analysis (OR 0.1, 95% CI 0.01 to 0.6), this finding was not statistically significant after IPW adjustments. In addition, those reporting possible exposure in the past 14 days had an OR of 5.0 (95% CI 1.0 to 25.1) in screening positive for depression, after adjusting for age, gender, smoking, customer-facing jobs, SARS-CoV-2 community prevalence in workers' residential cities and workers' self-reported history of anxiety and depression. The ability to practice social distancing consistently at work was inversely associated with both anxiety and depression, with adjusted OR 0.3 (95% CI 0.1 to 0.9) and 0.2 (95% CI 0.03 to 0.99), respectively. Moreover, those commuting to work by foot, bike or private car demonstrated a 90% risk reduction in screening positive for depression (OR 0.1, 95% CI 0.02 to 0.7) after accounting for potential confounders. In the sensitivity analysis using truncated IPW, all significant results remained robust.

In further sensitivity analysis, we categorised the workers' jobs into supervisory positions and non-supervisory positions. There were 7 out of 21 (33%) SARS-CoV-2-positive employees with supervisory positions, while among those tested negative for SARS-CoV-2 only 7.2% held a supervisory position ($p=0.005$). After using truncated IPW to adjust for age, gender, smoking and SARS-CoV-2 community prevalence, those with supervisory positions had an OR of 6.0 (95% CI 1.5 to 24.9) of having positive SARS-CoV-2 testing results.

Table 2 Characteristics of retail essential employees in a single grocery store in Massachusetts, USA presented for SARS-CoV-2, the virus causing COVID-19, RT-PCR assay testing by Generalised Anxiety Disorder 7-item scale (GAD-7) screening score for anxiety

	At least mild anxiety (GAD-7 >4) (N=24)	No or minimal anxiety (GAD-7 score ≤4) (N=75)	P value
Age, mean (SD)	45.5 (13.7)	50.0 (14.2)	0.169
Female, n (%)	15 (62.5%)	32 (42.7%)	0.145
Smoker, n (%)	6 (25.0%)	18 (24.0%)	0.999
Daily alcohol consumption, n (%)	2 (8.3%)	6 (8.0%)	0.999*
Marijuana use, n (%)	6 (25.0%)	7 (9.3%)	0.103
Self-reported exposure to SARS-CoV-2-positive individual(s) in the past 14 days, n (%)	9 (37.5%)	15 (20.0%)	0.142
Job positions with direct customer exposure at store†, n (%)	16 (66.7%)	48 (64.0%)	0.999
Full-time employment status, n (%)	19 (79.2%)	49 (65.3%)	0.308
Ability to practice social distancing consistently at work, count (%)	11 (45.8%)	57 (76.0%)	0.009
Using gloves consistently at work, count (%)	19 (79.2%)	58 (77.3%)	0.886
Wearing face mask consistently at work, count (%)	22 (91.7%)	70 (93.3%)	0.999*
Wearing face mask consistently outside of work, count (%)	15 (62.5%)	63 (84.0%)	0.072
Commute to work by foot, bike, or private car	18 (75.0%)	67 (89.3%)	0.156
PPE knowledge score, median (IQR)	15 (14–15)	15 (14–15)	0.867‡
COVID-19 perception score, median (IQR)	13 (11.5–15)	12 (11–14.75)	0.090‡

*Statistics derived from Fisher's exact test.

†Direct customer exposure positions include cashier, front end associate, sales associate, fresh food associate, cart attendant, janitorial crew, supervisor and manager of all levels. These are in contrast to positions mainly dealing with consumer goods or the environment, such as stocker, backroom, receiving and maintenance.

‡Statistics derived from Wilcoxon rank sum test with continuity correction.

PPE, personal protective equipment; RT-PCR, reverse transcriptase PCR.

Table 3 Characteristics of retail essential employees in a single grocery store in Massachusetts, USA presented for SARS-CoV-2, the virus causing COVID-19, RT-PCR assay testing by Patient Health Questionnaire-9 (PHQ-9) screening score for depression

	At least mild depression (PHQ-9 score >4) (N=8)	No or minimal depression (PHQ-9 score ≤4) (N=91)	P value
Age, mean (SD)	40.3 (10.5)	49.7 (14.2)	0.070
Female, n (%)	6 (75.0%)	41 (45.1%)	0.145*
Smoker, n (%)	3 (37.5%)	21 (23.1%)	0.397*
Daily alcohol consumption, n (%)	2 (25.0%)	6 (6.6%)	0.125*
Marijuana use, n (%)	2 (25.0%)	11 (12.1%)	0.282*
Self-reported exposure to SARS-CoV-2-positive individual(s) in the past 14 days, n (%)	5 (62.5%)	19 (20.9%)	0.019*
Job positions with direct customer exposure at store†, n (%)	6 (75.0%)	58 (63.7%)	0.712*
Full-time employment status, n (%)	7 (87.5%)	61 (67.0%)	0.429*
Ability to practice social distancing consistently at work, count (%)	2 (25.0%)	66 (72.5%)	0.010*
Using gloves consistently at work, count (%)	6 (75.0%)	71 (78.0%)	0.667*
Wearing face mask consistently at work, count (%)	7 (87.5%)	85 (93.4%)	0.409*
Wearing face mask consistently outside of work, count (%)	4 (50.0%)	74 (81.3%)	0.133*
Commuting to work by foot, bike or private car	4 (50.0%)	81 (89.0%)	0.013*
PPE knowledge score, median (IQR)	14.5 (14–15)	15 (14–15)	0.885‡
COVID-19 perception score, median (IQR)	13 (12–14)	12 (11–15)	0.402‡

*Statistics derived from Fisher's exact test.

†Direct customer exposure positions include cashier, front end associate, sales associate, fresh food associate, cart attendant, janitorial crew, supervisor and manager of all levels. These are in contrast to positions mainly dealing with consumer goods or the environment, such as stocker, backroom, receiving and maintenance.

‡Statistics derived from Wilcoxon rank sum test with continuity correction.

PPE, personal protective equipment; RT-PCR, reverse transcriptase PCR.

DISCUSSION

Our current study presents multiple valuable COVID-19-related associations in a group of essential workers during the pandemic. First, the infection rate of 20% positive SARS-CoV-2 RT-PCR assay results at this grocery retail store was significantly higher than the surrounding communities. In addition, most of these employees were asymptomatic at time of testing. After IPW adjustments, employees with direct exposure to customers had more than five times increased odds to have a positive SARS-CoV-2 RT-PCR assay result. We also found the ability to practice social distancing at workplace was inversely correlated to workers' anxiety and depression status. Lastly, having a confirmed SARS-CoV-2 exposure history in past 14 days and

commuting to work by public transportation or shared rides was strongly associated with depressive mood. To the best of our knowledge, this study is the first to report the above associations in a cohort of grocery retail essential employees.

There is limited research discussing non-HCWs essential workers in this pandemic, particularly retail employees and their exposure to customers.⁹ The SARS-CoV-2 infection rate among these retail employees was significantly higher than of the local community around similar time period, which was 0.9%–1.3%.²³ Previous studies on HCWs suggested COVID-19 infections among HCWs were consistent with community exposure rather than work-related exposure, with the prevalence ranging from 0% to 14%.^{13,14} In fact, a pioneering study conducted in the

Table 4 Crude and inverse probability weighting (IPW) adjusted ORs of positive SARS-CoV-2 (the virus causing COVID-19) RT-PCR assay, positive Generalised Anxiety Disorder 7-item scale (GAD-7) and positive Patient Health Questionnaire-9 (PHQ-9) screenings by key risk factors among retail essential employees in a single grocery store in Massachusetts, USA

	Crude OR (95% CI)	IPW adjusted OR (95% CI)	Truncated IPW adjusted OR (95% CI)
Positive SARS-CoV-2 RT-PCR assay			
Job positions with direct customer exposure at store*	6.2 (1.6 to 40.6)	5.1 (1.1 to 24.8)†	5.3 (1.1 to 25.6)†
Cigarette smoker	0.1 (0.01 to 0.6)	0.2 (0.02 to 1.4)‡	0.2 (0.02 to 1.3)‡
Positive GAD-7 screening (GAD-7 >4)			
Ability to practice social distancing consistently at work	0.3 (0.1 to 0.7)	0.3 (0.1 to 0.9)§	0.3 (0.1 to 0.8)§
Positive PHQ-9 screening (PHQ-9 score >4)			
Self-reported exposure to SARS-CoV-2-positive individual(s) in the past 14 days	6.3 (1.4 to 33.1)	5.0 (1.0 to 25.1)§	5.0 (1.0 to 24.9)§
Ability to practice social distancing consistently at work	0.1 (0.02 to 0.6)	0.2 (0.03 to 0.99)§	0.2 (0.03 to 0.9)§
Commute to work by foot, bike or private car	0.1 (0.03 to 0.6)	0.1 (0.02 to 0.7)§	0.1 (0.03 to 0.7)§

*Direct customer exposure positions include cashier, front end associate, sales associate, fresh food associate, cart attendant, janitorial crew, supervisor and manager of all levels. These are in contrast to positions mainly dealing with consumer goods or the environment, such as stocker, backroom, receiving and maintenance.

†IPW adjusting for age, gender, smoking and SARS-CoV-2 community prevalence in workers' residential cities.

‡IPW adjusting for age, gender, job positions with direct customer exposure, and SARS-CoV-2 community prevalence of workers' residential cities.

§IPW adjusting for age, gender, smoking, customer-facing jobs, SARS-CoV-2 community prevalence of workers' residential cities and self-reported history of anxiety and/or depression.

RT-PCR, reverse transcriptase PCR; Truncated IPW, IPW with extreme weights (below 5th and above 95th percentile) truncated.

Netherlands investigated the viral genetic sequences of affected HCWs and found the infection was more likely to be acquired from the communities.²⁴ In our current study, we did not observe a difference in SARS-CoV-2 community prevalence among those tested positive versus negative employees, indicating the possibility of a true work-related SARS-CoV-2 exposure. In terms of exposure risk, >90% of employees with positive assay result had a position with significant direct exposure to customers. We also found that employees in supervisory positions, with exposure from both customers and colleagues, had increased SARS-CoV-2 exposure risk. Employees in supervisory positions may have more exposure due to frequent interpersonal contacts, therefore leading to their higher infection rates. Notably, most of the SARS-CoV-2-positive assay workers were asymptomatic at time of testing. As evidence has shown probable transmission from asymptomatic or mildly symptomatic carriers,^{3 25 26} these workers as a cluster carries significant risk to their customers, colleagues and families. Our findings further strengthens the retail cluster transmission observed in a previous study from China, which involved supermarket employees, clients and the families of affected cases, resulting in a infection rate of 9.2% among the market workers.¹⁷

In this cohort, cigarette smoking was found to be a protective factor of SARS-CoV-2 RT-PCR assay result in the crude analysis. Despite a lack of statistical significance after IPW adjustment, our finding echoes a recently published systematic review indicating lower smoking prevalence among patients with COVID-19 in comparison with the general population.²⁷ In that review, the authors pooled 13 Chinese studies on hospitalised patients with COVID-19 and found a prevalence of 6.5% of current smokers, which was around one-fourth of the smoking prevalence among the general population. The potential biological mechanism involving nicotinic receptors has been proposed in another study.²⁸ In fact, research has shown nicotinic receptor activity can promote SARS-CoV-2 transmission through co-expression of ACE2 receptor, the host receptor for the virus. Therefore, the competitive nature of nicotine and SARS-CoV-2, as a nicotinic agent, for the receptor may serve as a key to prevent the infection.²⁸ Our finding of fewer current smokers with a positive SARS-CoV-2 assay result, while in agreement with recent epidemiological studies, contradicts common perception and clinical recommendation on risks and effects of cigarette smoking on lung health warranting further research investigations.²⁹

While previous research has raised concerns on psychological distress due to COVID-19 in addition to physiological threats on essential workers,¹¹ most of them were focused on HCWs.^{10 15 30–32} The prevalence of anxiety among HCWs in other countries ranged from 20% to 65% during the COVID-19 pandemic.^{15 30 32} In our study, 24% of these workers had at least mild anxiety, suggesting non-HCWs essential employees experience similar level of psychological distress. Contrary to common beliefs on the association between sufficient PPE and employees' psychological distress,^{33 34} the inability to practice social distancing consistently at work was a significant risk factor for anxiety and depression in this essential worker cohort. While we are unable to discern the direction of the effect due to the cross-sectional nature of this study, these mental health findings support the need to implement further preventive strategies and to provide additional mental health assistance to essential employees.

Our current study has several limitations. First, our limited sample size may prevent identification of certain associations that may require larger statistical power, and incidental findings may by chance be observed in a small sample-sized study.

However, the large effect sizes (ie, ORs) are unlikely to be entirely biased by unmeasured confounding factors. Second, this is a cross-sectional study and therefore causal relationship could not be inferred. At the same time, survey collection was conducted prior to SARS-CoV-2 RT-PCR sampling, suggesting our major findings should be free of reverse causation and any recall bias would be minimised. Third, while a majority of the employees from this retail store were tested at this designated location, some employees received testing at other clinics due to insurance, scheduling and/or location convenience. As this was a city-mandated testing, employees were assigned by the retail headquarter to be tested at this location if they had not received or scheduled to receive SARS-CoV-2 testing. Selection was neither based on their exposure risk nor health outcome and therefore the current study should be free of selection bias. Lastly, since our data collection was largely based on self-reported questionnaire, we incur unavoidable risk of measurement error, misclassification and related information bias.

At the same time, our study enjoys several strengths. First, the SARS-CoV-2 RT-PCR assay samples were collected by nasopharyngeal approach which provides the highest test sensitivity among all methods³⁵ and the outcomes of interest were assessed by validated screening tools including GAD-7 and PHQ-9. The possibility of outcome misclassification was therefore minimised. Second, our secondary sensitivity analysis results were in accordance with the main analysis which further strengthened our findings. Third, our study participants were restricted to grocery retail employees from one store and such restriction could eliminate potential confounding factors such as socioeconomic status. Lastly, we included all workers that were scheduled and presented to the testing tent during group testing days without any exclusion criteria. As a result of our strengths, findings in this study may be generalised to grocery store employees working during the COVID-19 pandemic in similar settings.

In conclusion, in this cohort of grocery retail essential workers, 20% had a positive SARS-CoV-2 RT-PCR assay result and the majority (76%) of them were asymptomatic at time of testing. Employees with direct customer exposure were five times more likely to have a positive SARS-CoV-2 assay result. The ability to social distance consistently at work was a significant protective factor for anxiety and depression. Commuting to work by public transportation/shared rides and having an exposure to a confirmed case within the past 14 days were positively associated with depression. Further research is warranted to investigate these associations and their public health implications among essential employees.

Contributors JY designed the study and collected the data. F-YL conducted data analysis and drafted the manuscript. F-YL, CS, SNK and JY all contributed to the interpretation of the data, revising the manuscript and final approval. JY supervised the project.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests SNK has received COVID-19-related consulting fees from Open Health.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement De-identified data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the

translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iD

Justin Yang <http://orcid.org/0000-0002-9743-2074>

REFERENCES

- Mahase E. Covid-19: WHO declares pandemic because of "alarming levels" of spread, severity, and inaction. *BMJ* 2020;368:m1036.
- Ghinai I, McPherson TD, Hunter JC, et al. First known person-to-person transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the USA. *Lancet* 2020;395:1137–44.
- Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med* 2020;382:970–1.
- Cheng H-Y, Jian S-W, Liu D-P, et al. Contact tracing assessment of COVID-19 transmission dynamics in Taiwan and risk at different exposure periods before and after symptom onset. *JAMA Intern Med* 2020. doi:10.1001/jamainternmed.2020.2020. [Epub ahead of print: 01 May 2020].
- Wang CJ, Ng CY, Brook RH. Response to COVID-19 in Taiwan: big data analytics, new technology, and proactive testing. *JAMA* 2020;323:1341–1342.
- Lai S, Ruktanonchai NW, Zhou L, et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature* 2020;585:410–3.
- Matrajt L, Leung T. Evaluating the effectiveness of social distancing interventions to delay or flatten the epidemic curve of coronavirus disease. *Emerg Infect Dis* 2020;26:1740–8.
- Sim MR. The COVID-19 pandemic: major risks to healthcare and other workers on the front line. *Occup Environ Med* 2020;77:281–2.
- Lan F-Y, Wei C-F, Hsu Y-T, et al. Work-Related COVID-19 transmission in six Asian countries/areas: a follow-up study. *PLoS One* 2020;15:e0233588.
- Kisely S, Warren N, McMahon L, et al. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* 2020;369:m1642.
- Gold JA. Covid-19: adverse mental health outcomes for healthcare workers. *BMJ* 2020;369:m1815.
- Wu P, Fang Y, Guan Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can J Psychiatry* 2009;54:302–11.
- Reusken CB, Buiting A, Bleeker-Rovers C, et al. Rapid assessment of regional SARS-CoV-2 community transmission through a convenience sample of healthcare workers, the Netherlands, March 2020. *Euro Surveill* 2020;25.
- Lan F-Y, Filler R, Mathew S, et al. COVID-19 symptoms predictive of healthcare workers' SARS-CoV-2 PCR results. *PLoS One* 2020;15:e0235460.
- Lai J, Ma S, Wang Y, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open* 2020;3:e203976.
- Wang L, Duan Y, Zhang W, et al. Epidemiologic and clinical characteristics of 26 cases of COVID-19 arising from patient-to-patient transmission in Liaocheng, China. *Clin Epidemiol* 2020;12:387–91.
- Zhang JZ, Zhou P, Han DB, et al. [Investigation on a cluster epidemic of COVID-19 in a supermarket in Liaocheng, Shandong province]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:E055.
- von Elm E, Altman DG, Egger M, et al. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335:806–8.
- Centers for Disease Control and Prevention (CDC). Interim guidelines for collecting, handling, and testing clinical specimens from persons for coronavirus disease, 2019. Available: <https://www.cdc.gov/coronavirus/2019-ncov/lab/guidelines-clinical-specimens.html> [Accessed 2 Sep 2020].
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–13.
- Spitzer RL, Kroenke K, Williams JBW, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166:1092–7.
- Pirracchio R, Resche-Rigon M, Chevret S. Evaluation of the propensity score methods for estimating marginal odds ratios in case of small sample size. *BMC Med Res Methodol* 2012;12:70.
- Massachusetts Department of Public Health. Massachusetts Department of Public Health COVID-19 Dashboard - Wednesday, 2020. Available: <https://www.mass.gov/doc/covid-19-dashboard-may-6-2020/download> [Accessed 2 Sep 2020].
- Sikkema RS, Pas SD, Nieuwenhuijse DF, et al. COVID-19 in health-care workers in three hospitals in the South of the Netherlands: a cross-sectional study. *Lancet Infect Dis* 2020;3099:30527–2.
- Kim ES, Chin BS, Kang CK, et al. Clinical course and outcomes of patients with severe acute respiratory syndrome coronavirus 2 infection: a preliminary report of the first 28 patients from the Korean cohort study on COVID-19. *J Korean Med Sci* 2020;35:e142.
- Bai Y, Yao L, Wei T, et al. Presumed asymptomatic carrier transmission of COVID-19. *JAMA* 2020;323:1406.
- Farsalinos K, Barbouni A, Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med* 2020.
- Lutchman D. Could the smoking gun in the fight against COVID-19 be the (rh)ACE-2? *Eur Respir J* 2020;56. doi:10.1183/13993003.01560-2020. [Epub ahead of print: 16 Jul 2020].
- World Health Organization. Who statement: tobacco use and COVID-19. Available: <https://www.who.int/news-room/detail/11-05-2020-who-statement-tobacco-use-and-covid-19> [Accessed 2 Sep 2020].
- Taghizadeh F, Hassannia L, Moosazadeh M, et al. Anxiety and depression in health workers and general population during COVID-19 epidemic in Iran: a web-based cross-sectional study. *medRxiv* 2020 2020.
- Spiller TR, Méan M, Ernst J, et al. Development of health care workers' mental health during the SARS-CoV-2 pandemic in Switzerland: two cross-sectional studies. *Psychol Med* 2020:1–4.
- Rossi R, Soggi V, Pacitti F, et al. Mental health outcomes among frontline and second-line health care workers during the coronavirus disease 2019 (COVID-19) pandemic in Italy. *JAMA Netw Open* 2020;3:e2010185.
- Kang L, Li Y, Hu S, et al. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry* 2020;7:e14.
- Nagesh S, Chakraborty S. Saving the frontline health workforce amidst the COVID-19 crisis: challenges and recommendations. *J Glob Health* 2020;10:010345.
- Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA* 2020.