

Occupational COVID-19: what can be learned from notifications of occupational diseases?

Marinaccio and colleagues are applauded for providing valuable data on the risk of acquiring COVID-19 due to exposure in the workplace in Italy.¹ Once again, this emphasises the importance of an assessment of the working circumstances and registration of occupational diseases. Data on work-related COVID-19 in the study of Marinaccio and colleagues was based on compensation claims of workers. In the Netherlands, an expert-based one-page guidance document for assessment and reporting of COVID-19 as occupational disease was developed by occupational physicians and data specialists. This guidance document, based on the general 6-step approach,² was published on the website of the Netherlands Center for Occupational Diseases³ and communicated by newsletters. Occupational COVID-19 reports, International Classification of Diseases-10 code U.071, contain information about age, sex, occupation, economic sector, work disability and potential cause for occupational COVID-19. In the period March–September 2020, 904 cases were reported.

Although this number is much lower compared to claims received in Italy reported by the Italian Workers' Compensation Authority, in the Netherlands, COVID-19 was the 2nd most common occupational disease reported from March–September (33% of all reported illnesses), exceed only by mental illness (42%). Musculoskeletal disorders accounted for 19%. This percentage is consistent with their estimate of 30% of reported occupational COVID-19 cases in the working age. Most Dutch occupational COVID-19 cases were reported within the nurse and caring homes (713; 79%) and hospitals (125; 14%), which is in line with the 70% reported for health-care workers in Ontario Canada.⁴ The other cases were mainly from the sectors 'wholesale and retail trade', 'administrative and support service activities', 'manufacturing', 'public administration and defence' and 'education'. Work disability

was reported in most of the cases and in 21% of the workers this lasted for more than 3 months. Among workers from nurse and caring homes, contact with infected patients and colleagues was reported in 94% of the cases. In 516 of the 670 cases (77%), unprotected contact with a patient with COVID-19 was reported, which could be caused by a lack of sufficient personal protective equipment. In 27 of the 670 cases (4%), nurses and caregivers were infected while using personal protective equipment.

Both approaches in assessment and reporting of occupational COVID-19 cases have own (dis)advantages. Compensation claim data may be biased by the financial need of a worker^{1 4} while the registry of occupational diseases may be biased by the reporting behaviour of occupational physicians.⁵ Nevertheless, notification and accompanying data are important to explore the impact of work-related risk factors and to identify groups at high risk of contracting an occupational disease. This is particularly relevant for new emerging occupational diseases such as COVID-19, when the knowledge on risk factors is limited but implementation of prevention measures is urged.

Recording of occupational diseases is of additional value in developing preventive measures in the workplace. To enable comparison of notification approaches used in different countries, harmonisation of epidemiological surveillance systems and case definitions such as currently done by the OMEGA-NET initiative⁶ is of the utmost importance.

Henk F van der Molen , **Sanja Kezic**, **Steven Visser**, **Gerda de Groene**, **Jaap Maas**, **Astrid de Wind**, **Sietske Tamminga** 

Amsterdam UMC, University of Amsterdam, Public and Occupational Health, Netherlands Center for Occupational Diseases, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands

Correspondence to Dr Henk F van der Molen, Public and Occupational Health, Amsterdam UMC Location AMC, Amsterdam 1105 AZ, The Netherlands; h.f.vandermolen@amsterdamumc.nl

Contributors All authors contributed to this letter.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.



To cite van der Molen HF, Kezic S, Visser S, *et al.* *Occup Environ Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/oemed-2020-107121

Received 5 October 2020

Revised 18 October 2020

Accepted 29 October 2020



► <http://dx.doi.org/10.1136/oemed-2020-106844>

Occup Environ Med 2020;0:1.

doi:10.1136/oemed-2020-107121

ORCID iDs

Henk F van der Molen <http://orcid.org/0000-0002-0719-2020>

Sietske Tamminga <http://orcid.org/0000-0002-2228-1187>

REFERENCES

- Marinaccio A, Bocconi F, Rondinone BM, *et al.* Occupational factors in the COVID-19 pandemic in Italy: compensation claims applications support establishing an occupational surveillance system. *Occup Environ Med* 2020. doi:10.1136/oemed-2020-106844. [Epub ahead of print: 23 Sep 2020].
- Boschman JS, Brand T, Frings-Dresen MHW, *et al.* Improving the assessment of occupational diseases by occupational physicians. *Occup Med* 2017;67:13–19.
- Netherlands Center for Occupational Diseases, 2020. Available: <https://www.beroepsziekten.nl/sites/default/files/documents/stappenplan-melden-beroepsziekte-covid-19-cascode-605.pdf>
- Institute for Work and Health, Canada, 2020. Available: <https://www.iwh.on.ca/newsletters/at-work/101/what-research-can-do-estimating-role-of-workplaces-in-covid-19-transmissions>
- van der Molen HF, Omvlee L, Brand T, *et al.* Perceived barriers and facilitators in the assessment of occupational diseases. *Occup Med* 2018;68:555–8.
- Turner MC, Mehlum IS. Greater coordination and harmonisation of European occupational cohorts is needed. *Occup Environ Med* 2018;75:475–6.