

**Supplemental Table 1: Frequency of autoimmune conditions by country**

	Overall	Australia	Israel	Spain	U.S.
	N=154	N=14	N=110	N=20	N=10
<b>Autoimmune disease findings, n (%)</b>					
Overall	33 (21.4)	7 (50.0)	22 (20.0)	1 (5.0)	3 (30.0)
Rheumatoid arthritis	9 (5.8)	0 (0.0)	6 (5.5)	0 (0.0)	3 (30.0)
Systemic sclerosis	7 (4.6)	0 (0.0)	7 (6.4)	0 (0.0)	0 (0.0)
Psoriasis/psoriatic arthritis	6 (3.9)	0 (0.0)	6 (5.5)	0 (0.0)	0 (0.0)
Abnormal autoimmune serology without a definitive autoimmune diagnosis	7 (4.5)	7 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other**	8 (5.2)	0 (0.0)	7 (6.4)	1 (5.0)	0 (0.0)

\*\*Other includes: Sjogren's syndrome, anti-neutrophil cytoplasmic autoantibody (ANCA) vasculitis, mixed connective tissue disease, cicatricial pemphigoid (rheumatic), systemic lupus erythematosus, or palindromic rheumatism.